

# On the Decoupling of the Homogeneous and Inhomogeneous Parts in Inhomogeneous Quantum Groups

Gaetano Fiore,

Dip. di Matematica e Applicazioni, Fac. di Ingegneria  
Università di Napoli, V. Claudio 21, 80125 Napoli

I.N.F.N., Sezione di Napoli,  
Complesso MSA, V. Cintia, 80126 Napoli

## Abstract

We show that, if there exists a realization of a Hopf algebra  $H$  in a  $H$ -module algebra  $\mathcal{A}$ , then one can split their cross-product into the tensor product algebra of  $\mathcal{A}$  itself with a subalgebra isomorphic to  $H$  and *commuting* with  $\mathcal{A}$ . This result applies in particular to the algebra underlying inhomogeneous quantum groups like the Euclidean ones, which are obtained as cross-products of the quantum Euclidean spaces  $\mathbb{R}_q^N$  with the quantum groups of rotation  $U_q so(N)$  of  $\mathbb{R}_q^N$ , for which it has no classical analog.

# 1 Introduction

As known, given a unital module algebra  $\mathcal{A}$  of a Lie algebra  $\mathfrak{g}$  (over the field  $\mathbb{C}$ , say), one can build a new module algebra, called cross-product  $U\mathfrak{g} \ltimes \mathcal{A}$ , that is as a vector space the tensor product  $\mathcal{A} \otimes U\mathfrak{g}$  of the vector spaces  $\mathcal{A}, U\mathfrak{g}$  (over the same field) and has the product law

$$(\mathbf{1}_{\mathcal{A}} \otimes g)(\mathbf{1}_{\mathcal{A}} \otimes g') = (\mathbf{1}_{\mathcal{A}} \otimes gg'), \quad (1.1)$$

$$(a \otimes \mathbf{1}_H)(a' \otimes \mathbf{1}_H) = aa' \otimes \mathbf{1}_H, \quad (1.2)$$

$$(a \otimes \mathbf{1}_H)(\mathbf{1}_{\mathcal{A}} \otimes g) = a \otimes g, \quad (1.3)$$

$$(\mathbf{1}_{\mathcal{A}} \otimes g)(a \otimes \mathbf{1}_H) = g_{(1)} \triangleright a \otimes g_{(2)}, \quad (1.4)$$

for any  $g, g' \in U\mathfrak{g}$ ,  $a, a' \in \mathcal{A}$ . Here we have denoted by  $\triangleright$  the left action of the Hopf algebra  $H \equiv U\mathfrak{g}$  on  $\mathcal{A}$ ,

$$\triangleright : (g, a) \in H \times \mathcal{A} \rightarrow g \triangleright a \in \mathcal{A}, \quad (1.5)$$

and used a Sweedler-type notation with suppressed summation sign for the coproduct  $\Delta(g)$  of  $g$ , namely the short-hand notation  $\Delta(g) = g_{(1)} \otimes g_{(2)}$  instead of a sum  $\Delta(g) = \sum_{\mu} g_{(1)}^{\mu} \otimes g_{(2)}^{\mu}$  of many terms. In the main part of this paper we shall work with a left action and therefore left-module algebras. In section 4 we shall give the formulae if we use instead a right action  $\triangleleft$  and right-module algebras. By definition of a (left) action, for any  $g, g' \in H$ ,  $a, a' \in \mathcal{A}$ ,

$$(gg') \triangleright a = g \triangleright (g' \triangleright a) \quad (1.6)$$

$$g \triangleright (aa') = (g_{(1)} \triangleright a)(g_{(2)} \triangleright a'); \quad (1.7)$$

We recall that the coproduct on the unit and on any  $g \in \mathfrak{g}$  is given by

$$\Delta(\mathbf{1}_H) = \mathbf{1}_H \otimes \mathbf{1}_H \quad \Delta(g) = g \otimes \mathbf{1}_H + \mathbf{1}_H \otimes g; \quad (1.8)$$

on the rest of  $U\mathfrak{g}$  it is determined using the fact that it is an algebra homomorphism  $\Delta : H \rightarrow H \otimes H$ . Clearly, the coproduct is cocommutative, i.e.  $g_{(1)} \otimes g_{(2)} = g_{(2)} \otimes g_{(1)}$ .

For the sake of clearness, above and in the sequel we denote the tensor products of vector spaces and of algebras by  $\otimes$  and  $\otimes$  (in boldface) respectively; therefore, given two unital algebras  $B, B'$  (over the same field),  $B \otimes B'$  is  $B \otimes B'$  as a vector space, while as an algebra it is characterized by the product

$$(a \otimes a')(b \otimes b') = (ab \otimes a'b') \quad (1.9)$$

for any  $a, b \in B$  and  $a', b' \in B'$ .

In the sequel, with a standard abuse of notation, for any unital algebras  $B, B'$  and any  $b \in B$ ,  $b' \in B'$  we shall denote by  $bb'$  the element  $b \otimes b'$  in the tensor product of vector spaces  $B \otimes B'$  and omit either unit  $\mathbf{1}_B, \mathbf{1}_{B'}$  whenever multiplied by non-unit elements (thus,  $\mathbf{1}_B B', B \mathbf{1}_{B'}$  will be denoted

by  $B', B$ ). Consequently, in the case of e.g. the cross product algebra  $U\mathfrak{g} \ltimes \mathcal{A}$  relations (1.1-1.3) take trivial forms, whereas (1.4) becomes the commutation relation

$$ga = (g_{(1)} \triangleright a)g_{(2)}; \quad (1.10)$$

whereas for a tensor product algebra  $B \otimes B'$  the analogs of relations (1.1-1.3) again take trivial forms, whereas (1.9) for  $a = \mathbf{1}_B$ ,  $b' = \mathbf{1}_{B'}$  becomes the trivial commutation relation

$$a' b = b a'. \quad (1.11)$$

Of course,  $B \otimes B'$  is isomorphic to  $B' \otimes B$ .

$H \ltimes \mathcal{A}$  is itself a module algebra under the left action  $\triangleright$  of  $H$  if we extend the latter on the elements of the  $H$  subalgebra as the adjoint action,

$$g \triangleright h = g_{(1)} h S g_{(2)} \quad g, h \in H; \quad (1.12)$$

(here  $S$  denotes the antipode of  $H$ ), and set as usual

$$g \triangleright (a \otimes h) = g_{(1)} \triangleright a \otimes g_{(2)} \triangleright h;$$

note that, in the notation mentioned above this relation takes the same form as (1.7), i.e. becomes  $g \triangleright (ah) = (g_{(1)} \triangleright a)(g_{(2)} \triangleright h)$ . It is immediate to show that relation (1.10) implies that one can realize the action  $\triangleright : H \times (H \ltimes \mathcal{A}) \rightarrow H \ltimes \mathcal{A}$  in the ‘adjoint-like way’

$$g \triangleright \eta = g_{(1)} \eta S g_{(2)} \quad (1.13)$$

on all of  $H \ltimes \mathcal{A}$ .

Classical examples of cross product algebras are the universal enveloping algebras of inhomogeneous Lie groups, like the Poincaré algebra, where  $H = U\mathfrak{so}(3,1)$  is the algebra generated by infinitesimal Lorentz transformations and  $\mathcal{A}$  is the abelian algebra generated by infinitesimal translations on Minkowski space, or the Euclidean algebra, where  $\mathcal{A}$  is the abelian algebra generated by infinitesimal translations on the  $N$ -dimensional Euclidean space  $\mathbb{R}^N$  and  $H = U\mathfrak{so}(N)$  is the algebra generated by its infinitesimal rotations.

The above setting can be “deformed” by allowing  $H$  to be a non-cocommutative Hopf algebra, e.g. the quantum group  $U_q \mathfrak{g}$ , and as  $\mathcal{A}$  the corresponding  $q$ -deformed module algebra. In general, the latter will be no more abelian, even if its classical counterpart is. A cross product  $H \ltimes \mathcal{A}$  can be still defined by means of the same formulae (1.5-1.10).

In this paper we want to show that there are prominent examples of cross products  $H \ltimes \mathcal{A}$  that are isomorphic to  $\mathcal{A} \otimes H$ , more precisely are equal to  $\mathcal{A} H'$  with  $H' \subset H \ltimes \mathcal{A}$  a subalgebra isomorphic to  $H$  and *commuting* with  $\mathcal{A}$ , even if this is not the case for their undeformed counterparts. As we shall see, this occurs if there exists an algebra homomorphism  $\varphi$  of the cross product into  $\mathcal{A}$  acting identically on the latter. Of course,

this will have dramatic consequences for the cross product both from the algebraic and from the representation-theoretic viewpoint; it will allow to reduce representations of  $H \ltimes \mathcal{A}$  to direct sums of tensor products of representations of  $\mathcal{A}$  and of  $H'$ . To prevent misunderstandings, we note that in general if  $H \ltimes \mathcal{A}$  itself is a Hopf algebra, as in the case of inhomogeneous quantum groups, neither  $\mathcal{A}$  nor  $H'$  will be a Hopf subalgebra of  $H \ltimes \mathcal{A}$ .

The present work has been inspired among others by the results of Ref. [3, 8]. In Ref. [3] the existence of such a  $H'$  for the  $q$ -deformed Euclidean algebra in three dimensions has been noted; its generators have been constructed “by hand” and have been used to decouple  $q$ -rotations from  $q$ -translations in the  $*$ -representations of  $H \ltimes \mathcal{A}$ . In Ref. [8] we had constructed “by hand” for the  $q$ -deformed Euclidean algebra in  $N \geq 3$  dimensions a set of generators which do the same job; but, instead of commuting with the  $q$ -deformed generators of translations, they  $q$ -commute with the latter. Now, *a posteriori*, one can check that they can be obtained as products of suitable elements of  $H'$  by suitable elements of the natural Cartan (i.e. maximal abelian) Hopf subalgebra of  $H = U_q so(N)$ .

On the contrary, the present work gives a very simple prescription for their construction, based on the existence of  $\varphi$ . The prescription can be thus applied to a number of models, including the following. In Ref. [2, 3] a class of homomorphisms (2.1-2.2) has been determined for a slightly enlarged version  $\mathcal{A}$  of the algebra of functions on the  $N$ -dimensional quantum Euclidean space  $\mathbb{R}_q^N$  or quantum Euclidean sphere  $S_q^{N-1}$ , the Hopf algebra  $H$  denoting  $U_q so(N)$  itself if  $N$  is odd, either the Borel subalgebra  $U_q^+ so(N)$  or the one  $U_q^- so(N)$  if  $N$  is even. Their behaviour under the  $*$ -structures has been investigated in Ref. [10] (where incidentally we draw another consequence of its existence, namely the possibility of “unbraiding” braided tensor product algebras). This will be explicitly described in section 5.1. The analogous maps for the  $q$ -deformed fuzzy sphere  $S_{q,M}^2$  have been found in [12]. On the other hand, the existence of algebra homomorphisms (2.1) for  $H = U_q so(N), U_q sl(N)$  and  $\mathcal{A}$  respectively equal to (a suitable completion of) the  $U_q so(N)$ -covariant Heisenberg algebra or the  $U_q sl(N)$ -covariant Heisenberg (or Clifford) algebras, has been known for even a longer time [7, 4, 13]. This will be treated in section 5.2. Note that in the latter cases  $\varphi$  exist also for the undeformed counterparts at  $q = 1$ , thus our results will apply also in this case (we do not know whether this has ever been formulated as a result in ordinary Lie group theory).

In section 2 we state and prove the main results of this work leading to the construction of  $H'$ . In section 3 we focus on the  $*$ -structures. In section 4 we give without proof for right-module algebras all the main formulae valid for left-module algebras. In section 5 we apply our results to the two examples of cross-product algebras mentioned above.

We conclude this section with some additional preliminaries. Beside the Sweedler-type notation with lower indices introduced for the coproduct, we shall denote a sum of many terms in a tensor product by a Sweedler-type

notation with *upper* indices and suppressed summation sign, e.g.  $c^{(1)} \otimes c^{(2)}$  will actually mean a sum  $\sum_{\mu} c_{\mu}^{(1)} \otimes c_{\mu}^{(2)}$ . Secondly, we can also introduce an ‘opposite’ action  $\triangleright^{op} : H \times (H \ltimes \mathcal{A}) \rightarrow H \ltimes \mathcal{A}$ , i.e. an action of the Hopf algebra with the same algebra structure and counit but opposite coproduct  $\Delta^{op}(g) = g_{(2)} \otimes g_{(1)}$  and inverse antipode, by

$$g \triangleright^{op} \eta = g_{(2)} \eta S^{-1} g_{(1)}. \quad (1.14)$$

It fulfills

$$(gg') \triangleright^{op} a = g \triangleright^{op} (g' \triangleright^{op} a) \quad (1.15)$$

$$g \triangleright^{op} (aa') = (g_{(2)} \triangleright^{op} a) (g_{(1)} \triangleright^{op} a'). \quad (1.16)$$

(Note that  $g \triangleright^{op} a$  in general is *not* an element of  $\mathcal{A}$ ).

## 2 The commutant of $\mathcal{A}$ within $H \ltimes \mathcal{A}$

### 2.1 The basic construction

In this subsection we assume that there exists an algebra homomorphism

$$\varphi : H \ltimes \mathcal{A} \rightarrow \mathcal{A} \quad (2.1)$$

acting as the identity on  $\mathcal{A}$ , namely for any  $a \in \mathcal{A}$

$$\varphi(a) = a. \quad (2.2)$$

(Note that, as a consequence,  $\varphi$  is idempotent:  $\varphi^2 = \varphi$ ). More explicitly, the fact that  $\varphi$  is a homomorphism implies that for any  $a \in \mathcal{A}$ ,  $g \in H$

$$\varphi(g)a = (g_{(1)} \triangleright a) \varphi(g_{(2)}). \quad (2.3)$$

[Note that (unless the left action of  $H$  on  $\mathcal{A}$  is trivial), no  $\varphi$  can exist if  $\mathcal{A}$  is abelian, e.g. for the Euclidean algebra  $Uso(N) \ltimes \mathbb{R}^N$ ; but, as we shall recall later,  $\varphi$  exists for the  $q$ -deformed Euclidean algebra, and therefore the result will apply.]

Let  $\mathcal{C}$  be the commutant of  $\mathcal{A}$  within  $H \ltimes \mathcal{A}$ , i.e. the subalgebra

$$\mathcal{C} := \{c \in H \ltimes \mathcal{A} \mid [c, a] = 0 \quad \forall a \in \mathcal{A}\}. \quad (2.4)$$

Clearly  $\mathcal{C}$  contains the center  $\mathcal{Z}(\mathcal{A})$  of  $\mathcal{A}$ . Let  $\zeta : H \rightarrow H \ltimes \mathcal{A}$  be the map defined by

$$\zeta(g) := \varphi(Sg_{(1)})g_{(2)}. \quad (2.5)$$

Note that if in the definition of  $\zeta$  we drop  $\varphi$ , we get instead the counit  $\varepsilon$ . Similarly, if we apply  $\varphi$  to  $\zeta$  and recall that  $\varphi$  is both a homomorphism and idempotent we also find

$$\varphi \circ \zeta = \varepsilon. \quad (2.6)$$

Now,  $\varepsilon(g)$  is a complex number times  $\mathbf{1}_{\mathcal{A}}$  and therefore trivially commutes with  $\mathcal{A}$ . Since  $\varphi$  does not change the commutation relations between  $\mathcal{A}$  and  $H$ , we expect that also  $\zeta(g)$  commutes with  $\mathcal{A}$ . This is confirmed by

**Theorem 1** *Let  $H$  be a Hopf algebra,  $\mathcal{A}$  a  $H$ -module algebra,  $\mathcal{C}$  the commutant (2.4), and  $\varphi$  an homomorphism of the type (2.1), (2.2). Then (2.5) defines an injective algebra homomorphism  $\zeta : H \rightarrow \mathcal{C}$ ; moreover  $\mathcal{C} = \mathcal{Z}(\mathcal{A})\zeta(H)$  and  $H \ltimes \mathcal{A} = \mathcal{A}\zeta(H)$ . If, in particular  $\mathcal{Z}(\mathcal{A}) = \mathbb{C}$ , then  $\mathcal{C} = \zeta(H)$  and  $\zeta : H \hookrightarrow \mathcal{C}$  is an algebra isomorphism.*

In other words, the subalgebra  $H'$  looked for in the introduction will be obtained by setting  $H' := \zeta(H)$ . For these reasons we shall call  $\zeta$ , as well as the other maps  $\zeta_i, \zeta_i^\pm$  which we shall introduce below, *decoupling maps*.

**Proof** For any  $a \in \mathcal{A}$ ,

$$\begin{aligned}
\zeta(g)a &\stackrel{(2.5)}{=} \varphi(Sg_{(1)})g_{(2)}a \\
&\stackrel{(1.10)}{=} \varphi(Sg_{(1)})(g_{(2)} \triangleright a)g_{(3)} \\
&\stackrel{(2.3)}{=} [Sg_{(2)} \triangleright (g_{(3)} \triangleright a)] \varphi(Sg_{(1)})g_{(4)} \\
&\stackrel{(1.6)}{=} [(Sg_{(2)}g_{(3)}) \triangleright a] \varphi(Sg_{(1)})g_{(4)} \\
&= [\varepsilon(g_{(2)})\mathbf{1}_H \triangleright a] \varphi(Sg_{(1)})g_{(3)} \\
&= a\varphi(Sg_{(1)})g_{(2)} = a\zeta(g),
\end{aligned} \tag{2.7}$$

proving that  $\zeta(g) \in \mathcal{C}$ . Using (2.7) with  $a = \varphi(Sg'_{(1)})$  we find

$$\begin{aligned}
\zeta(gg') &\stackrel{(2.5)}{=} \varphi[S(g_{(1)}g'_{(1)})]g_{(2)}g'_{(2)} = \varphi(Sg'_{(1)})\varphi(Sg_{(1)})g_{(2)}g'_{(2)} \\
&\stackrel{(2.5)}{=} \varphi(Sg'_{(1)})\zeta(g)g'_{(2)} \stackrel{(2.7)}{=} \zeta(g)\varphi(Sg'_{(1)})g'_{(2)} \\
&\stackrel{(2.5)}{=} \zeta(g)\zeta(g'),
\end{aligned} \tag{2.8}$$

proving that  $\zeta$  is a homomorphism. To prove that  $\zeta$  is injective note that  $\zeta(g) = \zeta(g')$  implies

$$\varphi(Sg_{(1)}) \otimes g_{(2)} = \varphi(Sg'_{(1)}) \otimes g'_{(2)},$$

whence, by applying  $(m \otimes \text{id}) \circ (\text{id} \otimes \varphi \otimes \text{id}) \circ (\text{id} \otimes \Delta)$  we find  $g = g'$  (we have denoted by  $m$  the multiplication map of  $\mathcal{A}$ ,  $m(a \otimes b) = ab$ ).

Now, consider a generic element  $c \in H \ltimes \mathcal{A}$  and decompose it in the form  $c = c^{(1)}c^{(2)}$  with  $c^{(1)} \otimes c^{(2)} \in \mathcal{A} \otimes H$ . From (2.5) it immediately follows that

$$c = c^{(1)}c^{(2)} = c^{(1)}\varphi\left(c^{(2)}_{(1)}\right)\zeta\left(c^{(2)}_{(2)}\right),$$

showing that  $H \ltimes \mathcal{A} = \mathcal{A}\zeta(H)$ , because  $c^{(1)}\varphi\left(c^{(2)}_{(1)}\right) \in \mathcal{A}$ . In particular, assume  $c \in \mathcal{C}$ . Then

$$0 = [a, c] = \left[a, c^{(1)}\varphi\left(c^{(2)}_{(1)}\right)\zeta\left(c^{(2)}_{(2)}\right)\right] \stackrel{(2.7)}{=} \left[a, c^{(1)}\varphi\left(c^{(2)}_{(1)}\right)\right]\zeta\left(c^{(2)}_{(2)}\right);$$

since  $\zeta$  is injective, all factors  $\zeta \left( c_{(2)}^{(2)} \right)$  are linearly independent and therefore

$$\left[ a, c^{(1)} \varphi \left( c_{(1)}^{(2)} \right) \right] = 0,$$

whence we conclude that  $c^{(1)} \varphi \left( c_{(1)}^{(2)} \right) \in \mathcal{Z}(\mathcal{A})$  and  $c \in \mathcal{Z}(\mathcal{A}) \zeta(H)$ .  $\square$

**Corollary 1** *Under the same assumptions of Thm. 1 the center of the cross-product  $H \ltimes \mathcal{A}$  is given by*

$$\mathcal{Z}(H \ltimes \mathcal{A}) = \mathcal{Z}(\mathcal{A}) \zeta(\mathcal{Z}(H)). \quad (2.9)$$

*Moreover, if  $H_c, \mathcal{A}_c$  are maximal abelian subalgebras of  $H$  and  $\mathcal{A}$  respectively, then  $\mathcal{A}_c \zeta(H_c)$  is a maximal abelian subalgebra of  $H \ltimes \mathcal{A}$ .*

This is almost all what we need in the determination of the Casimirs and of a complete set of commuting observables of a quantum system whose algebra of observables is equal to (or contains)  $H \ltimes \mathcal{A}$ , as in Ref. [8, 3]. In addition we just need that these two subalgebras be closed under the corresponding  $*$ -structure of  $H \ltimes \mathcal{A}$ , what will be investigated in section 3.

**Proof** The proof of the second statement is immediate. As for the first, if  $c \in \mathcal{Z}(H)$  then  $[c, H] = 0$  and, applying the homomorphism  $\zeta$ ,  $[\zeta(c), \zeta(H)] = 0$ ; on the other hand  $[\zeta(c), \mathcal{A}] = 0$  by (2.7). Hence,  $\zeta(c)$  commutes with  $\mathcal{A} \zeta(H)$ , i.e. with  $H \ltimes \mathcal{A}$ , by theorem 1. Similarly if  $\tilde{c} \in \mathcal{Z}(\mathcal{A})$  then  $[\tilde{c}, \mathcal{A}] = 0$ ; on the other hand  $[\tilde{c}, \zeta(H)] = 0$  by (2.7). Hence  $\tilde{c}$  commutes with  $H \ltimes \mathcal{A}$ . Therefore  $\mathcal{Z}(\mathcal{A}) \zeta(\mathcal{Z}(H)) \subset \mathcal{Z}(H \ltimes \mathcal{A})$ .

Viceversa, by theorem 1 any  $c \in H \ltimes \mathcal{A}$  can be expressed in the form  $c = c^{(1)} \zeta(c^{(2)})$  with  $c^{(1)} \otimes c^{(2)} \in \mathcal{A} \otimes H$ . If in particular  $c \in \mathcal{Z}(H \ltimes \mathcal{A})$ , then it must be on one hand

$$0 = [c^{(1)} \zeta(c^{(2)}), \mathcal{A}] = [c^{(1)}, \mathcal{A}] \zeta(c^{(2)}),$$

implying  $c^{(1)} \in \mathcal{Z}(\mathcal{A})$  by the linear independence of all factors  $\zeta(c^{(2)})$ ; on the other hand it must be

$$0 = [c^{(1)} \zeta(c^{(2)}), \zeta(H)] = c^{(1)} [\zeta(c^{(2)}), \zeta(H)] = c^{(1)} \zeta([c^{(2)}, H]),$$

implying  $c^{(2)} \in \mathcal{Z}(H)$ , by the linear independence of all factors  $c^{(1)}$  and the injectivity of  $\zeta$ . Therefore  $\mathcal{Z}(H \ltimes \mathcal{A}) \subset \mathcal{Z}(\mathcal{A}) \zeta(\mathcal{Z}(H))$ .  $\square$

Using the results of the theorem we easily show that the restrictions of the left action of  $H$  to  $\varphi(H)$  and  $H$  itself [see (1.12)] look the same:

**Proposition 1** *Under the same assumptions of Thm. 1, on the images of  $\varphi$  the left action reads*

$$g \triangleright \varphi(h) = \varphi(g \triangleright h); \quad (2.10)$$

*equivalently,*

$$g \varphi(h) = \varphi(g_{(1)} \triangleright h) g_{(2)}. \quad (2.11)$$

**Proof**

$$\begin{aligned}
\varphi(g \triangleright h) &\stackrel{(1.12)}{=} \varphi(g_{(1)} h S g_{(2)}) = \varphi(g_{(1)}) \varphi(h) \varphi(S g_{(2)}) \\
&\stackrel{(2.5)}{=} \varphi(g_{(1)}) \varphi(h) \zeta(g_{(2)}) S g_{(3)} \stackrel{(2.7)}{=} \varphi(g_{(1)}) \zeta(g_{(2)}) \varphi(h) S g_{(3)} \\
&\stackrel{(2.5)}{=} g_{(1)} \varphi(h) S g_{(2)} \stackrel{(1.12)}{=} g \triangleright \varphi(h).
\end{aligned}$$

□

Apart from  $\zeta_1 \equiv \zeta$ , other maps fulfilling the same property (2.6) are

$$\begin{aligned}
\zeta_2(g) &:= g_{(2)} \varphi(S^{-1} g_{(1)}) \\
\zeta_3(g) &:= S g_{(1)} \varphi(g_{(2)}) & \zeta_4(g) &:= \varphi(g_{(2)}) S^{-1} g_{(1)} \\
\zeta_5(g) &:= g_{(1)} \varphi(S g_{(2)}) & \zeta_6(g) &:= \varphi(S^{-1} g_{(2)}) g_{(1)} \\
\zeta_7(g) &:= S^{-1} g_{(2)} \varphi(g_{(1)}) & \zeta_8(g) &:= \varphi(g_{(1)}) S g_{(2)}
\end{aligned} \tag{2.12}$$

One could wonder whether they also fulfill the previous theorems. Using (2.11) one can easily show that:

- $\zeta_2 = \zeta$ ; (2.13)
- $\zeta_3 = \zeta \circ S$ ,  $\zeta_4 = \zeta_2 \circ S^{-1} = \zeta \circ S^{-1}$  so that  $\zeta_3(g), \zeta_4(g) \in \mathcal{C}$ , but  $\zeta_3, \zeta_4$  are antihomomorphisms;
- $\zeta_5, \zeta_6, \zeta_7, \zeta_8$  do *not* map  $H$  into  $\mathcal{C}$ .

Let us prove for instance the first statement:

$$\begin{aligned}
\zeta_2(g) &\stackrel{(2.12)}{=} g_{(2)} \varphi(S^{-1} g_{(1)}) \stackrel{(2.11)}{=} \varphi(g_{(2)} \triangleright S^{-1} g_{(1)}) g_{(3)} \\
&\stackrel{(1.12)}{=} \varphi(g_{(2)} S^{-1} g_{(1)} S g_{(3)}) g_{(4)} = \varphi(S g_{(1)}) g_{(2)} \stackrel{(2.5)}{=} \zeta(g).
\end{aligned}$$

How does  $\zeta(H)$  transform under the action  $\triangleright$  of  $H$ ? One can easily verify that it is not mapped into itself. On the contrary, under the opposite action  $\triangleright^{op}$  it is:

**Proposition 2** *Under the same assumptions of Thm. 1, for any  $g, h \in H$*

$$h \triangleright^{op} \zeta(g) = \zeta(h \triangleright^{op} g), \tag{2.14}$$

$$h \zeta(g) = \zeta(h_{(2)} \triangleright^{op} g) h_{(1)}. \tag{2.15}$$

**Proof** By (1.13), (2.15) is a direct consequence of (2.14). To prove the latter,

$$\begin{aligned}
\zeta(h \triangleright^{op} g) &\stackrel{(1.14)}{=} \zeta(h_{(2)} g S^{-1} h_{(1)}) \stackrel{(2.8)}{=} \zeta(h_{(2)}) \zeta(g) \zeta(S^{-1} h_{(1)}) \\
&\stackrel{(2.5), (2.13)}{=} \zeta_2(h_{(3)}) \zeta(g) \varphi(h_{(2)}) S^{-1} h_{(1)}
\end{aligned}$$



$$\begin{aligned}
&\stackrel{(2.7)}{=} \zeta_2(h_{(3)})\varphi(h_{(2)})\zeta(g)S^{-1}h_{(1)} \\
&\stackrel{(2.12)}{=} h_{(4)}\varphi(S^{-1}h_{(3)})\varphi(h_{(2)})\zeta(g)S^{-1}h_{(1)} \\
&= h_{(2)}\zeta(g)S^{-1}h_{(1)} \stackrel{(1.14)}{=} h \triangleright^{op} \zeta(g).
\end{aligned}$$

□

## 2.2 Construction adapted to Gauss decompositions of $H$

In view of the applications that we shall consider in section 5 it is now useful to consider the case that, instead of a  $\varphi$  we just have at our disposal two homomorphisms  $\varphi^+, \varphi^-$

$$\varphi^\pm : H^\pm \ltimes \mathcal{A} \rightarrow \mathcal{A} \quad (2.16)$$

fulfilling (2.2), where  $H^+, H^-$  denote two Hopf subalgebras of  $H$  such that Gauss decompositions  $H = H^+H^- = H^-H^+$  hold. (The typical case is when  $H = U_q\mathfrak{g}$  and  $H^+, H^-$  denote its positive and negative Borel subalgebras.) Then the theorems listed so far will apply separately to  $H^+ \ltimes \mathcal{A}$  and  $H^- \ltimes \mathcal{A}$ , if we define corresponding maps  $\zeta^\pm : H^\pm \rightarrow \mathcal{A}$  by

$$\zeta^\pm(g) := \varphi^\pm(Sg_{(1)})g_{(2)}, \quad (2.17)$$

where  $g \in H^\pm$  respectively. What can we say about the whole  $H \ltimes \mathcal{A}$ ? We now prove

**Theorem 2** *Let  $H$  be a Hopf algebra,  $\mathcal{A}$  a  $H$ -module algebra,  $\mathcal{C}$  the commutant (2.4), and  $\varphi^\pm$  homomorphisms of the type (2.16), (2.2). Under the above assumptions formulae (2.17) define injective algebra homomorphisms  $\zeta^\pm : H^\pm \rightarrow \mathcal{C}$ . Moreover,*

$$\mathcal{C} = \mathcal{Z}(\mathcal{A})\zeta^+(H^+)\zeta^-(H^-) = \mathcal{Z}(\mathcal{A})\zeta^-(H^-)\zeta^+(H^+) \quad (2.18)$$

and

$$H \ltimes \mathcal{A} = \mathcal{A}\zeta^+(H^+)\zeta^-(H^-) = \mathcal{A}\zeta^-(H^-)\zeta^+(H^+). \quad (2.19)$$

*In particular, if  $\mathcal{Z}(\mathcal{A}) = \mathbb{C}$ , then  $\mathcal{C} = \zeta^+(H^+)\zeta^-(H^-) = \zeta^-(H^-)\zeta^+(H^+)$ .*

**Proof** As anticipated, the fact that  $\zeta^\pm$  are injective algebra homomorphisms  $\zeta^\pm : H^\pm \rightarrow \mathcal{C}$  follows from theorem 1. Now, by the Gauss decomposition  $H = H^+H^-$  a generic element  $c \in H \ltimes \mathcal{A}$  can be decomposed in the form  $c = c^{(1)}c^{(2)}c^{(3)}$  with  $c^{(1)} \otimes c^{(2)} \otimes c^{(3)} \in \mathcal{A} \otimes H^+ \otimes H^-$ ; again, the Sweedler-type notation at the rhs means that a sum of many terms is understood. From (2.17) it immediately follows that

$$\begin{aligned}
c &= c^{(1)}c^{(2)}c^{(3)} \\
&= c^{(1)}\varphi^+\left(c_{(1)}^{(2)}\right)\zeta^+\left(c_{(2)}^{(2)}\right)\varphi^-\left(c_{(1')}^{(3)}\right)\zeta^-\left(c_{(2')}^{(3)}\right) \\
&\stackrel{(2.7)}{=} \zeta^+\left(c_{(2)}^{(2)}\right)\zeta^-\left(c_{(2')}^{(3)}\right)c^{(1)}\varphi^+\left(c_{(1)}^{(2)}\right)\varphi^-\left(c_{(1')}^{(3)}\right),
\end{aligned}$$

showing that  $H \ltimes \mathcal{A} = \mathcal{A} \zeta^+(H^+) \zeta^-(H^-)$ . In particular, assume  $c \in \mathcal{C}$ . Then

$$\begin{aligned} 0 &= [a, c] = \left[ a, c^{(1)} \varphi^+ \left( c_{(1)}^{(2)} \right) \varphi^- \left( c_{(1')}^{(3)} \right) \zeta^+ \left( c_{(2)}^{(2)} \right) \zeta^- \left( c_{(2')}^{(3)} \right) \right] \\ &\stackrel{(2.7)}{=} \left[ a, c^{(1)} \varphi^+ \left( c_{(1)}^{(2)} \right) \varphi^- \left( c_{(1')}^{(3)} \right) \right] \zeta^+ \left( c_{(2)}^{(2)} \right) \zeta^- \left( c_{(2')}^{(3)} \right); \end{aligned}$$

since  $\zeta^+, \zeta^-$  are injective, all factors  $\zeta^+ \left( c_{(2)}^{(2)} \right) \zeta^- \left( c_{(2')}^{(3)} \right)$  are linearly independent and therefore

$$\left[ a, c^{(1)} \varphi^+ \left( c_{(1)}^{(2)} \right) \varphi^- \left( c_{(1')}^{(3)} \right) \right] = 0,$$

whence we conclude that  $c^{(1)} \varphi^+ \left( c_{(1)}^{(2)} \right) \varphi^- \left( c_{(1')}^{(3)} \right) \in \mathcal{Z}(\mathcal{A})$  and  $c$  belongs to  $\mathcal{Z}(\mathcal{A}) \zeta^+(H^+) \zeta^-(H^-)$ .

The proof of the claims with  $\zeta^+(H^+), \zeta^-(H^-)$  in the inverse order follow in the same way from the Gauss decomposition  $H = H^- H^+$ .  $\square$

As a consequence of this theorem, for any  $g^+ \in H^+, g^- \in H^-$  there exists a sum  $c^{(1)} \otimes c^{(2)} \otimes c^{(3)} \in \mathcal{Z}(\mathcal{A}) \otimes H^- \otimes H^+$  (depending on  $g^+, g^-$ ) such that

$$\zeta^+(g^+) \zeta^-(g^-) = c^{(1)} \zeta^-(c^{(2)}) \zeta^+(c^{(3)}). \quad (2.20)$$

These will be the “commutation relations” between elements of  $\zeta^+(H^+)$  and  $\zeta^-(H^-)$ . Their form will depend on the specific algebras considered. In section 5 we shall determine these commutation relations for two examples of cross products with  $H = U_q \mathfrak{g}$  using the Faddeev-Reshetikhin-Takhtadjan generators of  $U_q \mathfrak{g}$ .

Of course propositions 1, 2 will still apply to the present situation if both  $g, h$  belong to  $H^+$  and we replace  $\varphi, \zeta$ , by  $\varphi^+, \zeta^+$  (or the same with  $+$  replaced by  $-$ ). What can we say otherwise?

**Proposition 3** *Under the same assumptions of Thm. 2.2, if  $g \in H^\pm, h \in H^\mp$*

$$g \triangleright \varphi^\mp(h) = \varphi^\pm(g_{(1)}) \varphi^\mp(h) \varphi^\pm(Sg_{(2)}), \quad (2.21)$$

$$g \varphi^\mp(h) = \varphi^\pm(g_{(1)}) \varphi^\mp(h) \varphi^\pm(Sg_{(2)}) g_{(3)}, \quad (2.22)$$

$$g \triangleright^{op} \zeta^\mp(h) = \zeta^\pm(g_{(2)}) \zeta^\mp(h) \zeta^\pm(S^{-1}g_{(1)}), \quad (2.23)$$

$$g \zeta^\mp(h) = \zeta^\pm(g_{(3)}) \zeta^\mp(h) \zeta^\pm(S^{-1}g_{(2)}) g_{(1)}. \quad (2.24)$$

The proof uses theorem 2.2 and is similar to the proofs of Propositions 1, 2. Similarly, the analog of Corollary 1 reads

**Corollary 2** *Under the same assumptions of Thm. 2.2, any element  $c$  of the center  $\mathcal{Z}(H \ltimes \mathcal{A})$  of the cross-product  $H \ltimes \mathcal{A}$  can be expressed in the form*

$$c = \zeta^+ \left( c^{(1)} \right) \zeta^- \left( c^{(2)} \right) c^{(3)}, \quad (2.25)$$

where  $c^{(1)} \otimes c^{(2)} \otimes c^{(3)} \in H^+ \otimes H^- \otimes \mathcal{Z}(\mathcal{A})$  and  $c^{(1)}c^{(2)} \otimes c^{(3)} \in \mathcal{Z}(H) \otimes \mathcal{Z}(\mathcal{A})$ ; viceversa any such object  $c$  is an element of  $\mathcal{Z}(H \bowtie \mathcal{A})$ . If  $H_c \subset H^+ \cap H^-$  and  $\mathcal{A}_c$  are maximal abelian subalgebras of  $H$  and  $\mathcal{A}$  respectively, then  $\mathcal{A}_c \zeta^+(H_c)$  (as well as  $\mathcal{A}_c \zeta^-(H_c)$ ) is a maximal abelian subalgebra of  $H \bowtie \mathcal{A}$ .

### 3 \*-structures

Assume that  $H$  is a Hopf  $*$ -algebra and  $\mathcal{A}$  a  $H$ -module  $*$ -algebra, which means that on  $H$  and  $\mathcal{A}$  there exist antilinear involutive antihomomorphisms, which we both denote by the symbol  $*$ , such that

$$[\varepsilon(g)]^* = \varepsilon(g^*) \quad (3.1)$$

$$(g^*)_{(1)} \otimes (g^*)_{(2)} = (g_{(1)})^* \otimes (g_{(2)})^* \quad (3.2)$$

$$(Sg)^* = S^{-1}g^* \quad (3.3)$$

$$(g \triangleright a_i)^* = (S^{-1}g^*) \triangleright a_i^* \quad (3.4)$$

[actually (3.3) is a consequence of (3.1), (3.2) and of the uniqueness of the antipode]. Then these two  $*$ -structures can be glued in a unique one to make  $H \bowtie \mathcal{A}$  a  $*$ -algebra itself. If

$$\varphi : H \bowtie \mathcal{A} \rightarrow \mathcal{A} \quad (3.5)$$

is a homomorphism acting as the identity on  $\mathcal{A}$ , then it is straightforward to check that  $\varphi' := * \circ \varphi \circ *$  also is. As a consequence, if in a concrete case we know that such a homomorphism is unique, then  $\varphi' = \varphi$  and we automatically conclude that it is a  $*$ -homomorphism,

$$\varphi(\alpha^*) = [\varphi(\alpha)]^*, \quad \alpha \in H \bowtie \mathcal{A}. \quad (3.6)$$

More generally, if the homomorphism  $\varphi$  is not unique, it is natural to look for one that is a  $*$ -homomorphism. We shall give explicit examples of this in section 5.

**Proposition 4** *If  $\varphi : H \bowtie \mathcal{A} \rightarrow \mathcal{A}$  is a  $*$ -homomorphism, then also the map  $\zeta : H \rightarrow \mathcal{C}$  is.*

**Proof**

$$\begin{aligned} \zeta(g^*) &\stackrel{(2.5)}{=} \varphi(Sg_{(1)}^*g_{(2)}^*) \stackrel{(3.3),(3.2)}{=} \varphi((S^{-1}g_{(1)})^*)g_{(2)}^* \\ &\stackrel{(3.6)}{=} [\varphi(S^{-1}g_{(1)})]^*g_{(2)}^* = [g_{(2)}\varphi(S^{-1}g_{(1)})]^* \\ &\stackrel{(2.12)}{=} [\zeta_2(g)]^* \stackrel{(2.13)}{=} [\zeta(g)]^*. \end{aligned}$$

□

Alternatively, it may happen that no  $*$ -homomorphism  $\varphi$  exists, but there exist homomorphisms  $\varphi^\pm$  of the type (2.16). If  $H^\pm$  are Hopf  $*$ -subalgebras (i.e. are closed under  $*$ ), then also  $\varphi'^\pm := * \circ \varphi^\pm \circ *$  are homomorphisms of the type (2.16), and as before we can look for  $\varphi^\pm$  that are  $*$ -homomorphisms,

$$\varphi^\pm(\alpha^*) = [\varphi^\pm(\alpha)]^*, \quad \alpha \in H^\pm \bowtie \mathcal{A}. \quad (3.7)$$

On the contrary, if  $H^\pm$  are mapped into each other by  $*$ , then  $* \circ \varphi^\pm \circ *$  is a homomorphism of the type  $\varphi^\mp$ , and we can look for one such that

$$\varphi^\pm(\alpha^*) = [\varphi^\mp(\alpha)]^*, \quad \alpha \in H^\mp \bowtie \mathcal{A}. \quad (3.8)$$

**Proposition 5** *If  $\varphi^\pm$  are  $*$ -homomorphism, then also the map  $\zeta^\pm : H^\pm \rightarrow \mathcal{C}$  are. If  $\varphi^\pm$  fulfill (3.8), then  $\zeta^\pm$  fulfill*

$$\zeta^\pm(g^*) = [\zeta^\mp(g)]^*, \quad g \in H^\mp. \quad (3.9)$$

**Proof** The first statement amounts to the preceding proposition applied to  $H^\pm \bowtie \mathcal{A}$ . As for the second, the proof is just a small variation:

$$\begin{aligned} \zeta^\pm(g^*) &\stackrel{(2.5)}{=} \varphi^\pm(Sg_{(1)}^*g_{(2)}^*) \stackrel{(3.3),(3.2)}{=} \varphi^\pm((S^{-1}g_{(1)})^*)g_{(2)}^* \\ &\stackrel{(3.8)}{=} [\varphi^\mp(S^{-1}g_{(1)})]^*g_{(2)}^* = [g_{(2)}\varphi^\mp(S^{-1}g_{(1)})]^* \\ &\stackrel{(2.12)}{=} [\zeta_2^\mp(g)]^* \stackrel{(2.13)}{=} [\zeta^\mp(g)]^*. \end{aligned}$$

□

## 4 Formulae for right-module algebras

In this section we give the analogs for right  $H$ -module algebras of the main results found so far for left  $H$ -module algebras. By definition the right action  $\triangleleft : \mathcal{A} \times H \rightarrow \mathcal{A}$  fulfills

$$a \triangleleft (gg') = (a \triangleleft g) \triangleleft g' \quad (4.1)$$

$$(aa') \triangleleft g = (a \triangleleft g_{(1)}) (a' \triangleleft g_{(2)}). \quad (4.2)$$

The algebra  $\mathcal{A} \bowtie H$  is defined as follows. As a vector space it is  $H \otimes \mathcal{A}$ , whereas the product is defined through formulae obtained from (1.1-1.3) by flipping the tensor factors, together with

$$(\mathbf{1}_H \otimes a)(g \otimes \mathbf{1}_\mathcal{A}) = g_{(1)} \otimes (a \triangleleft g_{(2)})$$

for any  $a \in \mathcal{A}$ ,  $g \in H$ . As before, we shall denote  $g \otimes a$  by  $ga$  and omit either unit  $\mathbf{1}_\mathcal{A}$ ,  $\mathbf{1}_H$  whenever multiplied by non-unit elements; consequently only the last condition takes a non-trivial form and becomes

$$ag = g_{(1)} (a \triangleleft g_{(2)}). \quad (4.3)$$

$\mathcal{A} \rtimes H$  itself is a  $H$ -module algebra under the right action  $\triangleleft$  of  $H$  if we extend the latter on the elements of  $H$  as the adjoint action,

$$h \triangleleft g = Sg_{(1)}hg_{(2)}, \quad g, h \in H; \quad (4.4)$$

Let  $\tilde{\mathcal{C}}$  be the commutant of  $\mathcal{A}$  within  $\mathcal{A} \rtimes H$ . Clearly it contains the center  $\mathcal{Z}(\mathcal{A})$  of  $\mathcal{A}$ . We shall need homomorphisms

$$\tilde{\varphi} : \mathcal{A} \rtimes H \rightarrow \mathcal{A} \quad (4.5)$$

acting as the identity on  $\mathcal{A}$ , namely for any  $a \in \mathcal{A}$

$$\tilde{\varphi}(a) = a. \quad (4.6)$$

**Theorem 3** *Let  $H$  be a Hopf algebra,  $\mathcal{A}$  a right  $H$ -module algebra,  $\tilde{\mathcal{C}}$  the commutant of  $\mathcal{A}$  within  $\mathcal{A} \rtimes H$ , and  $\tilde{\varphi}$  an homomorphism of the type (4.5), (4.6). Then the map  $\zeta_5$  defined by (2.12) (with  $\varphi$  replaced by  $\tilde{\varphi}$ ) is an injective algebra homomorphism  $\zeta_5 : H \rightarrow \tilde{\mathcal{C}}$ ; moreover  $\tilde{\mathcal{C}} = \zeta_5(H) \mathcal{Z}(\mathcal{A})$  and  $\mathcal{A} \rtimes H = \zeta_5(H) \mathcal{A}$ . If, in particular  $\mathcal{Z}(\mathcal{A}) = \mathbb{C}$ , then  $\tilde{\mathcal{C}} = \zeta(H)$  and  $\zeta_5 : H \hookrightarrow \tilde{\mathcal{C}}$  is an algebra isomorphism.*

Consider the maps  $\zeta_i$  defined by (2.12), but with  $\varphi$  replaced by  $\tilde{\varphi}$ . One can easily show that:

•

$$\zeta_6 = \zeta_5; \quad (4.7)$$

- $\zeta_8 = \zeta_6 \circ S = \zeta_5 \circ S$ ,  $\zeta_7 = \zeta_5 \circ S^{-1}$ , so that  $\zeta_7(g), \zeta_8(g) \in \mathcal{C}$ , but  $\zeta_7, \zeta_8$  are antihomomorphisms;
- $\zeta_1, \zeta_2, \zeta_3, \zeta_4$  do not map  $H$  into  $\tilde{\mathcal{C}}$ .

**Proposition 6** *If  $g, h \in H$*

$$\tilde{\varphi}(h) \triangleleft g = \tilde{\varphi}(h \triangleleft g), \quad (4.8)$$

$$\tilde{\varphi}(h)g = g_{(1)}\tilde{\varphi}(h \triangleleft g_{(2)}). \quad (4.9)$$

If, instead of a  $\tilde{\varphi}$ , we just have at our disposal two homomorphisms  $\tilde{\varphi}^+, \tilde{\varphi}^-$

$$\tilde{\varphi}^\pm : \mathcal{A} \rtimes H^\pm \rightarrow \mathcal{A}, \quad (4.10)$$

where  $H^+, H^-$  denote two Hopf subalgebras of  $H$  such that Gauss decompositions  $H = H^+H^- = H^-H^+$  hold, then the theorems listed so far will apply separately to  $\mathcal{A} \rtimes H^+$  and  $\mathcal{A} \rtimes H^-$ , if we define corresponding maps  $\zeta_5^\pm : H^\pm \rightarrow \mathcal{A}$  by

$$\zeta_5^\pm(g) := g_{(1)}\tilde{\varphi}^\pm(Sg_{(2)}), \quad (4.11)$$

where  $g \in H^\pm$  respectively. More precisely:

**Theorem 4** *Under the above assumptions formulae (4.11) define injective algebra homomorphisms  $\zeta_5^\pm : H^\pm \rightarrow \tilde{\mathcal{C}}$ . Moreover,*

$$\tilde{\mathcal{C}} = \zeta_5^+(H^+) \zeta_5^-(H^-) \mathcal{Z}(\mathcal{A}) = \zeta_5^-(H^-) \zeta_5^+(H^+) \mathcal{Z}(\mathcal{A}) \quad (4.12)$$

and

$$\mathcal{A} \rtimes H = \zeta_5^+(H^+) \zeta_5^-(H^-) \mathcal{A} = \zeta_5^-(H^-) \zeta_5^+(H^+) \mathcal{A}. \quad (4.13)$$

In particular, if  $\mathcal{Z}(\mathcal{A}) = \mathbb{C}$ , then  $\tilde{\mathcal{C}} = \zeta_5^+(H^+) \zeta_5^-(H^-) = \zeta_5^-(H^-) \zeta_5^+(H^+)$ .

**Proposition 7** *If  $g \in H^\pm$ ,  $h \in H^\mp$*

$$\tilde{\varphi}^\mp(h) \triangleleft g = \tilde{\varphi}^\pm(Sg_{(1)}) \tilde{\varphi}^\mp(h) \tilde{\varphi}^\pm(g_{(2)}), \quad (4.14)$$

$$\tilde{\varphi}^\mp(h) g = g_{(1)} \tilde{\varphi}^\pm(Sg_{(2)}) \tilde{\varphi}^\mp(h) \tilde{\varphi}^\pm(g_{(3)}). \quad (4.15)$$

If  $H$  is a Hopf  $*$ -algebra and  $\mathcal{A}$  a  $H$ -module  $*$ -algebra, then the two  $*$ -structures of  $H$  and  $\mathcal{A}$  can be glued into a unique one to make  $\mathcal{A} \rtimes H$  a  $*$ -algebra itself.

**Proposition 8** *If  $\tilde{\varphi} : \mathcal{A} \rtimes H \rightarrow \mathcal{A}$  is a  $*$ -homomorphism, then also the map  $\zeta_5 : H \rightarrow \tilde{\mathcal{C}}$  is.*

**Proposition 9** *If the maps  $\tilde{\varphi}^\pm$  defined in (4.11) are  $*$ -homomorphism, then also the maps  $\zeta_5^\pm : H^\pm \rightarrow \mathcal{C}$  are. If  $\tilde{\varphi}^\pm$  fulfill*

$$\tilde{\varphi}^\pm(\alpha^*) = [\tilde{\varphi}^\mp(\alpha)]^*, \quad \alpha \in \mathcal{A} \rtimes H^\mp, \quad (4.16)$$

then  $\zeta_5^\pm$  fulfill

$$\zeta_5^\pm(g^*) = [\zeta_5^\mp(g)]^*, \quad g \in H^\mp. \quad (4.17)$$

## 5 Applications

We now consider a couple of applications where  $H$  is the quantum group  $U_q \mathfrak{g}$  [5], with  $\mathfrak{g} = sl(N)$  or  $\mathfrak{g} = so(N)$ . As a set of generators of  $U_q \mathfrak{g}$  it is convenient to introduce the FRT generators [6]  $\mathcal{L}_j^{+i}, \mathcal{L}_j^{-i}$  ( $i, j$  take  $N$  different values), together with the square roots of the diagonal elements  $\mathcal{L}_j^{+j}, \mathcal{L}_j^{-j}$ . In the appendix we recall the relations they fulfill. The FRT generators are related to the so-called universal  $R$ -matrix  $\mathcal{R}$  by

$$\mathcal{L}_l^{+a} := \mathcal{R}^{(1)} \rho_l^a(\mathcal{R}^{(2)}) \quad \mathcal{L}_l^{-a} := \rho_l^a(\mathcal{R}^{-1(1)}) \mathcal{R}^{-1(2)}, \quad (5.1)$$

where we have denoted by  $\rho$  the fundamental  $N$ -dimensional representation of  $U_q sl(N)$  or  $U_q so(N)$ . Since in our conventions  $\mathcal{R} \in H^+ \otimes H^-$  ( $H^+, H^-$  denote the positive, negative Borel subalgebras) we see that  $\mathcal{L}_l^{+a} \in H^+$  and  $\mathcal{L}_l^{-a} \in H^-$ .

For historical reasons we introduce algebras  $\mathcal{A}$  as right- (rather than left-)  $U_q \mathfrak{g}$ -module algebras.

## 5.1 The Euclidean quantum group $\mathbb{R}_q^N \rtimes U_q so(N)$

As algebra  $\mathcal{A}$  we shall consider a slight extension of the quantum Euclidean space  $\mathbb{R}_q^N$  [6] (the  $U_q so(N)$ -covariant quantum space), i.e. of the unital associative algebra generated by  $p^i$  fulfilling the relations

$$\mathcal{P}_a^{ij} p^h p^k = 0, \quad (5.2)$$

where  $\mathcal{P}_a$  denotes the  $q$ -deformed antisymmetric projector appearing in the decomposition of the braid matrix  $\hat{R}$  of  $U_q so(N)$  [given in formula (A.10)]; the latter is related to  $\mathcal{R}$  by  $\hat{R}_{hk}^{ij} = \rho_h^j(\mathcal{L}_k^{+i}) = (\rho_h^j \otimes \rho_k^i)(\mathcal{R})$ . The multiplet  $(p^i)$  carries the fundamental  $N$ -dim (or vector) representation  $\rho$  of  $U_q so(N)$ : for any  $g \in U_q so(N)$

$$p^i \triangleleft g = \rho_j^i(g) p^j. \quad (5.3)$$

This implies

$$p^i \mathcal{L}_b^{\pm a} = \mathcal{L}_c^{\pm a} p^j \hat{R}_{jb}^{\pm 1 ci}, \quad (5.4)$$

$$S \mathcal{L}_b^{\pm a} p^i = \hat{R}^{\pm 1 ai}_{jk} p^j S \mathcal{L}_b^{\pm k}. \quad (5.5)$$

To define  $\tilde{\varphi}$  or  $\tilde{\varphi}^\pm$  one [2] slightly enlarges  $\mathbb{R}_q^N$  as follows. One introduces some new generators  $\sqrt{P_a}$ , with  $1 \leq a \leq \frac{N}{2}$ , together with their inverses  $(\sqrt{P_a})^{-1}$ , requiring that

$$P_a^2 = \sum_{h=-a}^a p^h p_h = \sum_{h,k=-a}^a g_{hk} p^h p^k. \quad (5.6)$$

In the previous equation  $g_{hk}$  denotes the ‘metric matrix’ of  $SO_q(N)$ :

$$g_{ij} = g^{ij} = q^{-\rho_i} \delta_{i,-j}. \quad (5.7)$$

It is a  $SO_q(N)$ -isotropic tensor and is a deformation of the ordinary Euclidean metric. Here and in the sequel  $n := \lfloor \frac{N}{2} \rfloor$  is the rank of  $so(N)$ , the indices take the values  $i = -n, \dots, -1, 0, 1, \dots, n$  for  $N$  odd, and  $i = -n, \dots, -1, 1, \dots, n$  for  $N$  even. We have also introduced the notation  $(\rho_i) = (n - \frac{1}{2}, \dots, \frac{1}{2}, 0, -\frac{1}{2}, \dots, \frac{1}{2} - n)$  for  $N$  odd,  $(n - 1, \dots, 0, 0, \dots, 1 - n)$  for  $N$  even. In the sequel we shall call  $P_n^2$  also  $P^2$ . Moreover for odd  $N$  we add also  $\sqrt{p^0}$  and its inverse as new generators. The commutation relations involving these new generators can be fixed consistently, and turn out to be simply  $q$ -commutation relations.  $P$  plays the role of ‘deformed Euclidean distance’ of the generic ‘point of coordinates’  $(p^i)$  of  $\mathbb{R}_q^N$  from the ‘origin’;  $P_a$  is the ‘projection’ of  $P$  on the ‘subspace’  $p^i = 0$ ,  $|i| > a$ . The center of  $\mathbb{R}_q^N$  is generated by  $\sqrt{P}$  and, only in the case of even  $N$ , by  $\sqrt{\frac{p^1}{p^{-1}}}$  and its inverse  $\sqrt{\frac{p^{-1}}{p^1}}$ . In the case of even  $N$  one needs to include also the FRT generator  $\mathcal{L}_1^{-1} = \mathcal{L}^{+1}_{-1}$  and its inverse  $\mathcal{L}_1^{+1} = \mathcal{L}^{-1}_{-1}$  [which are generators of  $U_q so(N)$  belonging to the natural Cartan subalgebra]

among the generators of  $\mathcal{A}$ . From (5.4) one derives that they satisfy the commutation relations

$$\mathcal{L}_1^{-1} p^{\pm 1} = q^{\pm 1} p^{\pm 1} \mathcal{L}_1^{-1}, \quad \mathcal{L}_1^{-1} p^{\pm i} = p^{\pm i} \mathcal{L}_1^{-1} \quad \text{for } i > 1 \quad (5.8)$$

with the generators of  $\mathcal{A}$ , and the standard FRT relations with the rest of  $U_q so(N)$ . As a consequence,  $\sqrt{\frac{p^{\pm 1}}{p^{\mp 1}}}$  are eliminated from the center of  $\mathcal{A}$  (in fact  $\mathcal{L}_1^{\pm 1}$  do not  $q$ -commute with  $\sqrt{\frac{p^{\pm 1}}{p^{\mp 1}}}$ ).

One can easily show that the extension of the action of  $U_q so(N)$  to  $\sqrt{P_a}, (\sqrt{P_a})^{-1}$  is uniquely determined by the constraints the latter fulfill; it is a bit complicated and therefore will be omitted, since we will not need its explicit expression. We keep the action of  $H$  on  $\mathcal{L}_1^{-1}$  as the standard (right) adjoint action (4.4). Note that the maps  $\tilde{\varphi}^{\pm}$  have no analog in the “undeformed” case ( $q = 1$ ), because  $\mathcal{A}_1 \equiv \mathbb{R}^N$  is abelian, whereas  $H \equiv U_q so(N)$  is not.

The homomorphisms [2]  $\tilde{\varphi}^{\pm} : \mathcal{A} \rtimes U_q^{\pm} so(N) \rightarrow \mathcal{A}$  take the simplest and most compact expression on the FRT generators of  $U_q^{\pm} so(N)$ . Let us introduce the short-hand notation  $[A, B]_x = AB - xBA$ . The images of  $\tilde{\varphi}^{-}$  on the negative FRT generators read

$$\tilde{\varphi}^{-}(\mathcal{L}_j^{-i}) = g^{ih} [\mu_h, p^k]_q g_{kj}, \quad (5.9)$$

where

$$\begin{aligned} \mu_0 &= \gamma_0 (p^0)^{-1} && \text{for } N \text{ odd,} \\ \mu_{\pm 1} &= \gamma_{\pm 1} (p^{\pm 1})^{-1} \mathcal{L}_1^{\pm 1} && \text{for } N \text{ even,} \\ \mu_a &= \gamma_a P_{|a|}^{-1} P_{|a|-1}^{-1} p^{-a} && \text{otherwise,} \end{aligned} \quad (5.10)$$

and  $\gamma_a \in \mathbb{C}$  are normalization constants fulfilling the conditions

$$\begin{aligned} \gamma_0 &= -q^{-\frac{1}{2}} h^{-1} && \text{for } N \text{ odd,} \\ \gamma_{\pm 1} &= -k^{-1} && \text{for } N \text{ even,} \\ \gamma_1 \gamma_{-1} &= -q^{-1} h^{-2} && \text{for } N \text{ odd,} \\ \gamma_a \gamma_{-a} &= -q^{-1} k^{-2} \omega_a \omega_{a-1} && \text{for } a > 1. \end{aligned} \quad (5.11)$$

Here  $h := q^{\frac{1}{2}} - q^{-\frac{1}{2}}$ ,  $k := q - q^{-1}$ ,  $\omega_a := (q^{\rho_a} + q^{-\rho_a})$ . On the other hand, the images of  $\tilde{\varphi}^{+}$  on the positive FRT generators read

$$\tilde{\varphi}^{+}(\mathcal{L}_j^{+i}) = g^{ih} [\bar{\mu}_h, p^k]_{q^{-1}} g_{kj}, \quad (5.12)$$

where

$$\begin{aligned} \bar{\mu}_0 &= \bar{\gamma}_0 (p^0)^{-1} && \text{for } N \text{ odd,} \\ \bar{\mu}_{\pm 1} &= \bar{\gamma}_{\pm 1} (p^{\pm 1})^{-1} \mathcal{L}_1^{\mp 1} && \text{for } N \text{ even,} \\ \bar{\mu}_a &= \bar{\gamma}_a P_{|a|}^{-1} P_{|a|-1}^{-1} p^{-a} && \text{otherwise,} \end{aligned} \quad (5.13)$$



and  $\bar{\gamma}_a \in \mathbb{C}$  normalization constants fulfilling the conditions

$$\begin{aligned}\bar{\gamma}_0 &= q^{\frac{1}{2}} h^{-1} && \text{for } N \text{ odd,} \\ \bar{\gamma}_{\pm 1} &= k^{-1} && \text{for } N \text{ even,} \\ \bar{\gamma}_1 \bar{\gamma}_{-1} &= -q h^{-2} && \text{for } N \text{ odd,} \\ \bar{\gamma}_a \bar{\gamma}_{-a} &= -q k^{-2} \omega_a \omega_{a-1} && \text{for } a > 1.\end{aligned}\tag{5.14}$$

From definition (2.12), using (A.13), (5.9), (5.12), (A.14), we find

$$\begin{aligned}\zeta_5^-(\mathcal{L}_j^{-i}) &= \mathcal{L}_h^{-i} \tilde{\varphi}^-(S\mathcal{L}_j^{-h}) = \mathcal{L}_h^{-i} [\mu_h, p^i]_q, \\ \zeta_5^+(\mathcal{L}_j^{+i}) &= \mathcal{L}_h^{+i} \tilde{\varphi}^+(S\mathcal{L}_j^{+h}) = \mathcal{L}_h^{+i} [\bar{\mu}_h, p^i]_{q^{-1}}.\end{aligned}\tag{5.15}$$

The case of even  $N$  is slightly outside the scheme developed in the preceding sections. As anticipated, the generators  $\mathcal{L}_1^{\pm 1}$  and their inverses  $\mathcal{L}_{-1}^{\pm 1}$  cannot be realized as (rational) ‘functions’ of the  $p$ ’s and have to be introduced in the codomain of  $\tilde{\varphi}^{\pm}$  as new generators. In fact definitions (5.9), (5.12) are exactly designed to lead to

$$\tilde{\varphi}^{\pm}(\mathcal{L}_1^{\pm 1}) = \mathcal{L}_1^{\pm 1} \quad \tilde{\varphi}^{\pm}(\mathcal{L}_{-1}^{\pm 1}) = \mathcal{L}_{-1}^{\pm 1}.\tag{5.16}$$

As a consequence, when  $s = \pm 1$

$$\zeta_5^+(\mathcal{L}_s^{+s}) = \mathcal{L}_k^{+s} \tilde{\varphi}^+(S\mathcal{L}_s^{+k}) \stackrel{(A.1), (5.16)}{=} \mathcal{L}_s^{+s} S \mathcal{L}_s^{+s} = 1\tag{5.17}$$

$$\zeta_5^-(\mathcal{L}_s^{-s}) = \mathcal{L}_k^{-s} \tilde{\varphi}^-(S\mathcal{L}_s^{-k}) \stackrel{(A.2), (5.16)}{=} \mathcal{L}_s^{-s} S \mathcal{L}_s^{-s} = 1.\tag{5.18}$$

Moreover, it is easy to check that

$$\begin{aligned}\mathcal{L}_1^{-1} \zeta_5^{\pm}(\mathcal{L}_j^{\pm i}) &= \zeta_5^{\pm}(\mathcal{L}_j^{\pm i}) \mathcal{L}_1^{-1} q^{\eta_i - \eta_j} \\ \mathcal{L}_1^{+1} \zeta_5^{\pm}(\mathcal{L}_j^{\pm i}) &= \zeta_5^{\pm}(\mathcal{L}_j^{\pm i}) \mathcal{L}_1^{+1} q^{\eta_j - \eta_i},\end{aligned}\tag{5.19}$$

where we have introduced the shorthand notation

$$\eta_i := \delta_i^1 - \delta_i^{-1}.\tag{5.20}$$

Therefore,  $\zeta_5^{\pm}[U_q^{\pm} so(2n)]$  do not commute with the whole  $\mathcal{A}$  but only with  $\mathbb{R}_q^{2n}$ , and the decomposition (4.12) will have to be modified into

$$\tilde{\mathcal{C}} = \zeta_5^+(H^+) \zeta_5^-(H^-) \mathcal{Z}(\mathbb{R}_q^{2n}) = \zeta_5^-(H^-) \zeta_5^+(H^+) \mathcal{Z}(\mathbb{R}_q^{2n})\tag{5.21}$$

with  $\tilde{\mathcal{C}}$  defined as the commutant of  $\mathbb{R}_q^{2n}$  within  $\mathbb{R}_q^{2n} \rtimes U_q so(2n)$ . As said,  $\mathcal{Z}(\mathbb{R}_q^{2n})$  contains also  $\sqrt{\frac{p^{\pm 1}}{p^{\mp 1}}}$ . Since  $\mu_{\pm 1}$ ,  $\bar{\mu}_{\mp 1}$  and therefore  $\tilde{\varphi}^-(\mathcal{L}_{\pm 1}^{-i})$ ,  $\tilde{\varphi}^+(\mathcal{L}_{\mp 1}^{+i})$  are of the form “ $\mathcal{L}_1^{\mp 1} \times$  an expression depending only on the  $p$ ’s”, they  $q$ -commute rather than commute with  $\zeta^{\pm}(\mathcal{L}_j^{\pm i})$ .

For odd and even  $N$  the commutation relations among the  $\zeta_5^-(\mathcal{L}_j^{-i})$  or among the  $\zeta_5^+(\mathcal{L}_j^{+i})$  are immediately obtained from (A.1-A.6) and (A.8) applying  $\zeta_5^-, \zeta_5^+$  and using the fact that they are homomorphisms.

To derive the commutation relations between the  $\zeta_5^-(\mathcal{L}^{-i}_j)$  and the  $\zeta_5^+(\mathcal{L}^{+h}_k)$  [the analog of (2.20) in explicit form] we need the commutation relations between the  $\tilde{\varphi}^-(\mathcal{L}^{-i}_j)$  and the  $\tilde{\varphi}^+(\mathcal{L}^{+i}_j)$ . To proceed we have to distinguish the case of odd and even  $N$ . In the formulation of the following lemma and proposition we switch off Einstein summation convention. In the appendix we prove

**Lemma 1** *If  $N$  is odd, then*

$$\tilde{\varphi}^-(S\mathcal{L}^{-i}_k)\tilde{\varphi}^+(S\mathcal{L}^{+h}_j) = \frac{\gamma_k}{\bar{\gamma}_k} \sum_{l,m,r,s} \hat{R}^{-1ih}_{lm} \tilde{\varphi}^+(S\mathcal{L}^{+l}_r) \tilde{\varphi}^-(S\mathcal{L}^{-m}_s) \hat{R}^{rs}_{kj} \frac{\bar{\gamma}_s}{\gamma_s}. \quad (5.22)$$

*If  $N$  is even, then if  $k \notin \{-1, 1\}$*

$$\begin{aligned} \tilde{\varphi}^-(S\mathcal{L}^{-i}_k)\tilde{\varphi}^+(S\mathcal{L}^{+h}_j) &= \frac{\gamma_k}{\bar{\gamma}_k} \sum_{l,m} \hat{R}^{-1ih}_{lm} \left[ \sum_{\substack{r,s \\ s \neq \pm 1}} \tilde{\varphi}^+(S\mathcal{L}^{+l}_r) \tilde{\varphi}^-(S\mathcal{L}^{-m}_s) \hat{R}^{rs}_{kj} \frac{\bar{\gamma}_s}{\gamma_s} \right. \\ &\quad \left. + \sum_{\substack{r,s \\ s = \pm 1}} \tilde{\varphi}^+(S\mathcal{L}^{+l}_r) \tilde{\varphi}^-(S\mathcal{L}^{-m}_{-s}) \frac{p^{-s}}{p^s} \hat{R}^{rs}_{kj} \frac{\bar{\gamma}_s}{\gamma_s} q^2 \right], \end{aligned} \quad (5.23)$$

*and if  $k \in \{-1, 1\}$*

$$\begin{aligned} \tilde{\varphi}^-(S\mathcal{L}^{-i}_k)\tilde{\varphi}^+(S\mathcal{L}^{+h}_j) &= - \sum_{l,m} \hat{R}^{-1ih}_{lm} \left[ \sum_{\substack{r,s \\ s \neq \pm 1}} \tilde{\varphi}^+(S\mathcal{L}^{+l}_r) \tilde{\varphi}^-(S\mathcal{L}^{-m}_s) \hat{R}^{rs}_{-k,j} \frac{\bar{\gamma}_s}{\gamma_s} \right. \\ &\quad \left. + \sum_{\substack{r,s \\ s = \pm 1}} \tilde{\varphi}^+(S\mathcal{L}^{+l}_r) \tilde{\varphi}^-(S\mathcal{L}^{-m}_{-s}) \frac{p^{-s}}{p^s} \hat{R}^{rs}_{-k,j} \frac{\bar{\gamma}_s}{\gamma_s} q^2 \right] \frac{p^{-k}}{p^k} q^{-2+2\eta_j\eta_k}. \end{aligned} \quad (5.24)$$

Note that for odd  $N$  if the ratio  $\frac{\gamma_a}{\bar{\gamma}_a}$  is independent of  $a$  (therefore it must be equal to  $\frac{\gamma_0}{\bar{\gamma}_0} = -q^{-1}$ ), the  $\gamma, \bar{\gamma}$ 's disappear, and by comparison with (A.7) we find [2] that setting  $\tilde{\varphi}(\mathcal{L}^{+i}_j) = \tilde{\varphi}^+(\mathcal{L}^{+i}_j)$ ,  $\tilde{\varphi}(\mathcal{L}^{-i}_j) = \tilde{\varphi}^-(\mathcal{L}^{-i}_j)$  defines a homomorphism  $\tilde{\varphi} : \mathcal{A} \rtimes U_{qso}(N) \rightarrow \mathcal{A}$ . For  $|q| = 1$   $\tilde{\varphi}$  turns out to be a  $*$ -homomorphism w.r.t. the corresponding  $*$ -structures (see below).

**Proposition 10** *If  $N$  is odd*

$$\zeta_5^+(\mathcal{L}^{+h}_j)\zeta_5^-(\mathcal{L}^{-i}_k) = \sum_{c,d,r,s} \hat{R}^{-1ih}_{cd} \zeta_5^-(\mathcal{L}^{-d}_s) \zeta_5^+(\mathcal{L}^{+c}_r) \hat{R}^{rs}_{kj} \frac{\gamma_k}{\bar{\gamma}_k} \frac{\bar{\gamma}_s}{\gamma_s}. \quad (5.25)$$

*If  $N$  is even, then*

$$\begin{aligned} \zeta_5^+(\mathcal{L}^{+h}_j)\zeta_5^-(\mathcal{L}^{-i}_k) &= \frac{\gamma_k}{\bar{\gamma}_k} q^{\eta_j(\eta_i - \eta_k)} \sum_{l,m,r} \hat{R}^{-1ih}_{lm} \left[ \sum_{s \neq \pm 1} \zeta_5^-(\mathcal{L}^{-m}_s) \zeta_5^+(\mathcal{L}^{+l}_r) \right. \\ &\quad \left. + \sum_{s = \pm 1} \zeta_5^-(\mathcal{L}^{-m}_{-s}) \zeta_5^+(\mathcal{L}^{+l}_r) \frac{p^{-s}}{p^s} q^{2+\eta_s(\eta_r - \eta_l)} \right] \hat{R}^{rs}_{kj} \frac{\bar{\gamma}_s}{\gamma_s} \end{aligned} \quad (5.26)$$

if  $k \notin \{1, -1\}$ , and, if  $k \in \{1, -1\}$ ,

$$\begin{aligned} \zeta_5^+(\mathcal{L}^{+h}_j) \zeta_5^-(\mathcal{L}^{-i}_k) &= -q^{-2+\eta_j(\eta_i+\eta_k)} \sum_{l,m,r} \hat{R}^{-1ih}_{lm} \left[ \sum_{s \neq \pm 1} \zeta_5^-(\mathcal{L}^{-m}_s) \zeta_5^+(\mathcal{L}^{+l}_r) \right. \\ &\quad \left. + \sum_{s=\pm 1} \zeta_5^-(\mathcal{L}^{-m}_{-s}) \zeta_5^+(\mathcal{L}^{+l}_r) \frac{p^{-s}}{p^s} q^{2+\eta_s(\eta_r-\eta_l)} \right] \zeta_5^+(\mathcal{L}^{+l}_r) \hat{R}^{rs}_{-k,j} \frac{\bar{\gamma}_s p^{-k}}{\gamma_s p^k}. \end{aligned} \quad (5.27)$$

So for even  $N$  the coefficients in the commutations relations depend explicitly on the central (in  $\mathbb{R}_q^N$ ) elements  $\frac{p^{\pm 1}}{p^{\mp 1}}$ .

Let us analyze now the properties of  $\zeta_5^\pm$  under  $*$ -structures. When  $|q| = 1$  the  $*$ -structure of  $\mathbb{R}_q^N$  is given by  $(p^i)^* = p^i$ . It turns out [10] that  $\tilde{\varphi}^\pm$  are  $*$ -homomorphisms if, in addition,

$$\gamma_a^* = -\gamma_a \begin{cases} 1 & \text{if } a < -1, \\ q^{-2} & \text{if } a > 1, \end{cases} \quad \begin{array}{l} \text{or } a = -1 \text{ and } N \text{ odd;} \\ \text{or } a = 1 \text{ and } N \text{ odd.} \end{array} \quad (5.28)$$

Under these assumptions, in view also of (A.15), we can apply proposition 9 and we conclude that  $\zeta_5^\pm$  are  $*$ -homomorphisms. When  $q \in \mathbb{R}^+$  the real structure of  $\mathbb{R}_q^N$  is given by  $(p^i)^* = p^j g_{ji}$ . It turns out [10] that  $\tilde{\varphi}^\pm$  fulfill  $[\varphi^\pm(g)]^* = \varphi^\mp(g^*)$ , or more explicitly

$$[\varphi^-(\mathcal{L}^{-i}_j)]^* = \varphi^+[(\mathcal{L}^{-i}_j)^*], \quad (5.29)$$

if, in addition,

$$\gamma_a^* = -\bar{\gamma}_{-a} \begin{cases} 1 & \text{if } a < -1, \\ q^{-2} & \text{if } a > 1, \end{cases} \quad \begin{array}{l} \text{or } a = -1 \text{ and } N \text{ odd;} \\ \text{or } a = 1 \text{ and } N \text{ odd.} \end{array} \quad (5.30)$$

Under these assumptions, in view also of (A.17), (A.15), we can apply proposition 9 and we conclude that  $\zeta_5^\pm$  fulfill

$$\zeta_5^\pm(g^*) = [\zeta_5^\mp(g)]^*, \quad g \in H^\mp. \quad (5.31)$$

## 5.2 The cross product of $U_q \mathfrak{g}$ 's with $U_q \mathfrak{g}$ -covariant Heisenberg algebras

As algebra  $\mathcal{A}$  we shall consider a slight extension of the  $U_q \mathfrak{g}$  -covariant deformed Heisenberg algebras  $\mathcal{D}_\epsilon \mathfrak{g}$ ,  $\mathfrak{g} = sl(N), so(N)$ . Such algebras have been introduced in Ref. [15, 16, 1]. They are unital associative algebras generated by  $x^i, \partial_j$  fulfilling the relations

$$\mathcal{P}_{a_{hk}}^{ij} x^h x^k = 0, \quad \mathcal{P}_{a_{hk}}^{ij} \partial_j \partial_i = 0, \quad \partial_i x^j = \delta_j^i + (q\gamma \hat{R})_{ih}^{\epsilon jk} x^h \partial_k, \quad (5.32)$$

where  $\gamma = q^{\frac{1}{N}}, 1$  respectively for  $\mathfrak{g} = sl(N), so(N)$ , and the exponent  $\epsilon$  can take either value  $\epsilon = 1, -1$ .  $\hat{R}$  denotes the braid matrix of  $U_q \mathfrak{g}$  [given in

formulae (A.9), (A.10)], and the matrix  $\mathcal{P}_a$  is the deformed antisymmetric projector appearing in the decompositions (A.11), (A.12) of the latter. The coordinates  $x^i$ , as the  $p^i$  of subsection 5.1, transform according to the fundamental  $N$ -dimensional representation  $\rho$  of  $U_q\mathbf{g}$ , whereas the ‘partial derivatives’ transform according the contragradient representation,

$$x^i \triangleleft g = \rho_j^i(g)x^j, \quad \partial_i \triangleleft g = \partial_h \rho_i^h(S^{-1}g). \quad (5.33)$$

In our conventions the indices will take the values  $i = 1, \dots, N$  if  $\mathbf{g} = sl(N)$ , whereas if  $\mathbf{g} = so(N)$  they will take the same values considered in subsection 5.1. In fact the quantum Euclidean space can be considered as a subalgebra of the  $U_qso(N)$ -covariant Heisenberg algebra, either by the identification  $p^i \equiv x^i$  or by the one  $p^i \equiv g^{ij}\partial_j$ .

Algebra homomorphisms  $\tilde{\varphi} : \mathcal{A} \rtimes H \rightarrow \mathcal{A}$ , for  $H = U_q\mathbf{g}$  and  $\mathcal{A}$  equal to (a suitable completion of)  $\mathcal{D}_{\epsilon, \mathbf{g}}$  have been constructed in Ref. [7, 4]. This is the  $q$ -analog of the well-known fact that the elements of  $\mathbf{g}$  can be realized as “vector fields” (first order differential operators) on the corresponding  $\mathbf{g}$ -covariant (undeformed) space, e.g.  $\tilde{\varphi}(E_j^i) = x^i\partial_j - \frac{1}{N}\delta_j^i$  in the  $\mathbf{g} = sl(N)$  case. This means that our decoupling map  $\zeta_5$  exists also in the undeformed case; we don’t know whether this result in Lie group representation theory has ever been formulated before.

The explicit expression of  $\tilde{\varphi}(\mathcal{L}_j^{-i})$  in terms of  $x^i, \partial_j$  for  $U_qsl(2), U_qso(3)$  has been given in Ref. [10]. For different values of  $N$  it can be found from the results of Ref. [7, 4] by passing from the generators adopted there to the FRT generators. For example, for  $\mathbf{g} = sl(2)$  and  $\epsilon = 1$  one finds

$$\begin{aligned} \tilde{\varphi}(\mathcal{L}_1^{+1}) &= \tilde{\varphi}(\mathcal{L}_2^{-2}) = [\tilde{\varphi}(\mathcal{L}_1^{-1})]^{-1} = [\tilde{\varphi}(\mathcal{L}_2^{+2})]^{-1} = \alpha\Lambda^{\frac{1}{2}} [1 + (q^2 - 1)x^2\partial_2]^{\frac{1}{2}} \\ \tilde{\varphi}(\mathcal{L}_2^{+1}) &= -\alpha k q^{-1} \Lambda^{\frac{1}{2}} [1 + (q^2 - 1)x^2\partial_2]^{-\frac{1}{2}} x^1\partial_2 \\ \tilde{\varphi}(\mathcal{L}_1^{-2}) &= \alpha k q^3 \Lambda^{\frac{1}{2}} [1 + (q^2 - 1)x^2\partial_2]^{-\frac{1}{2}} x^2\partial_1, \end{aligned} \quad (5.34)$$

where  $\alpha$  is fixed by (A.7) to be  $\alpha = \pm 1, \pm i$  and we have set

$$\Lambda^{-2} := 1 + (q^2 - 1)x^i\partial_i. \quad (5.35)$$

Whereas for  $\mathbf{g} = so(3)$  and  $\epsilon = 1$  one finds on the positive Borel subalgebra

$$\begin{aligned} \tilde{\varphi}(\mathcal{L}^{+-}) &= -\alpha\Lambda [1 + (q - 1)x^0\partial_0 + (q^2 - 1)x^+\partial_+] \\ \tilde{\varphi}(\mathcal{L}_0^{+-}) &= \alpha k \Lambda (x^-\partial_0 - \sqrt{q}x^0\partial_+) \\ \tilde{\varphi}(\mathcal{L}_+^{+-}) &= \frac{1}{1+q^{-1}}\tilde{\varphi}(\mathcal{L}_0^{+-})\tilde{\varphi}(\mathcal{L}_+^{0+}) \\ \tilde{\varphi}(\mathcal{L}_0^{0+}) &= 1 \\ \tilde{\varphi}(\mathcal{L}_+^{0+}) &= -q^{-\frac{1}{2}}[\tilde{\varphi}(\mathcal{L}^{+-})]^{-1}\tilde{\varphi}(\mathcal{L}_0^{+-}) \\ \tilde{\varphi}(\mathcal{L}_+^{++}) &= [\tilde{\varphi}(\mathcal{L}^{+-})]^{-1} \end{aligned} \quad (5.36)$$

and on the negative Borel subalgebra

$$\begin{aligned}
\tilde{\varphi}(\mathcal{L}^{--}) &= -(\alpha\Lambda[1 + (q-1)x^0\partial_0 + (q^2-1)x^+\partial_+])^{-1} \\
\tilde{\varphi}(\mathcal{L}^{-0}_-) &= -\alpha q^2 k \tilde{\varphi}(\mathcal{L}^{--}) \Lambda(x^0\partial_- - \sqrt{q}x^+\partial_0) \\
\tilde{\varphi}(\mathcal{L}^{-+}_-) &= \frac{1}{1+q} \tilde{\varphi}(\mathcal{L}^{-+}_0) \tilde{\varphi}(\mathcal{L}^{-0}_-) \\
\tilde{\varphi}(\mathcal{L}^{-0}_0) &= 1 \\
\tilde{\varphi}(\mathcal{L}^{-+}_0) &= -\alpha q^{\frac{3}{2}} k \Lambda(x^0\partial_- - \sqrt{q}x^+\partial_0) \\
\tilde{\varphi}(\mathcal{L}^{-+}_+) &= [\tilde{\varphi}(\mathcal{L}^{--})]^{-1}.
\end{aligned} \tag{5.37}$$

Here we have set

$$\Lambda^{-2} := [1 + (q^2 - 1)x^i\partial_i + \frac{(q^2 - 1)^2}{\omega_1^2}(g_{ij}x^ix^j)(g^{hk}\partial_k\partial_h)], \tag{5.38}$$

where

$$\omega_a := (q^{\rho_a} + q^{-\rho_a}),$$

and replaced for simplicity the values  $-1, 0, 1$  of the indices by the ones  $-, 0, +$ . In either case the  $\tilde{\varphi}$ -images of  $\mathcal{L}^{+i}_j$  and  $\mathcal{L}^{-j}_i$  for  $i > j$  vanish, because the latter do.

We see that strictly speaking  $\tilde{\varphi}$  takes values in some appropriate completion of  $\mathcal{D}_{\epsilon, \mathbf{g}}$ , containing at least the square root and inverse square root of the polynomial  $\Lambda^{-2}$  respectively defined in (5.35), (5.38), as well as the square root of  $[1 + (q^2 - 1)x^2\partial_2]$  and its inverse, when  $\mathbf{g} = sl(2)$ , and the inverses (5.36)<sub>6</sub>, (5.37)<sub>6</sub>, when  $\mathbf{g} = so(3)$ . Apart from this minimal completion, another possible one is the so-called  $h$ -adic, namely the ring of formal power series in  $h = \log q$  with coefficients in  $\mathcal{D}_{\epsilon, \mathbf{g}}$ . Other completions, e.g. in operator norms, can be considered according to the needs. One can easily show that the extension of the action of  $H$  to any such completion is uniquely determined (we omit to write down its explicit expression, since we don't need it).

$\mathcal{A} \rtimes H$  is a  $*$ -algebra and the map  $\tilde{\varphi}$  is a  $*$ -homomorphism both for  $q$  real and  $|q| = 1$ . The  $*$ -structure of  $\mathcal{A}$  is

$$(x^i)^* = x^i, \quad (\partial_i)^* = -\partial_i \begin{cases} q^{\pm 2(N-i+1)} & \text{if } H = U_q sl(N) \\ q^{\pm N + \rho_i} & \text{if } H = U_q so(N) \end{cases} \tag{5.39}$$

if  $|q| = 1$ , and

$$(x^h)^* = x^k g_{kh}, \quad (\partial_i)^* = -\frac{\Lambda^{\pm 2}}{q^{\pm N} + q^{\pm 2}} \left[ (g^{jh}\partial_h\partial_j), x^i \right] \tag{5.40}$$

if  $H = U_q so(N)$  and  $q \in \mathbb{R}^+$ . The upper or lower sign respectively refer to the choices  $\epsilon = 1, -1$  in (5.32)<sub>3</sub>, and  $\Lambda^{\pm 2}$  are respectively defined by

$$\Lambda^{\pm 2} := \left[ 1 + (q^{\pm 2} - 1)x^i\partial_i + \frac{(q^{\pm 2} - 1)^2}{\omega_n^2} r^2 (g^{ji}\partial_i\partial_j) \right]^{-1}. \tag{5.41}$$

In either case, in view also of (A.17), we can apply proposition 8 and we conclude that  $\zeta_5$  is a  $*$ -homomorphism.

## A Appendix

### Basic properties of $U_q\mathfrak{g}$

For both  $H = U_qsl(N), U_qso(N)$  the FRT generators fulfill the relations

$$\mathcal{L}_j^{+i} = 0, \quad \text{if } i > j, \quad (\text{A.1})$$

$$\mathcal{L}_j^{-i} = 0, \quad \text{if } i < j, \quad (\text{A.2})$$

$$\mathcal{L}_i^{-i}\mathcal{L}_i^{+i} = \mathcal{L}_i^{+i}\mathcal{L}_i^{-i} = 1, \quad \forall i \quad (\text{A.3})$$

$$\prod_i \mathcal{L}_i^{+i} = 1, \quad \prod_i \mathcal{L}_i^{-i} = 1, \quad (\text{A.4})$$

$$\hat{R}_{cd}^{ab} \mathcal{L}_f^d \mathcal{L}_e^+ = \mathcal{L}_c^+ \mathcal{L}_d^a \hat{R}_{ef}^{dc}, \quad (\text{A.5})$$

$$\hat{R}_{cd}^{ab} \mathcal{L}_f^d \mathcal{L}_e^- = \mathcal{L}_c^- \mathcal{L}_d^a \hat{R}_{ef}^{dc}, \quad (\text{A.6})$$

$$\hat{R}_{cd}^{ab} \mathcal{L}_f^d \mathcal{L}_e^- = \mathcal{L}_c^- \mathcal{L}_d^a \hat{R}_{ef}^{dc}; \quad (\text{A.7})$$

in addition, when  $H = U_qso(N)$  they also fulfill

$$\mathcal{L}_j^{\pm i} \mathcal{L}_k^{\pm h} g^{kj} = g^{hi} \quad \mathcal{L}_i^{\pm j} \mathcal{L}_h^{\pm k} g_{kj} = g_{hi}. \quad (\text{A.8})$$

Here  $\hat{R}_{cd}^{ab}$  denotes the braid matrix of  $U_qsl(N), U_qso(N)$ , and  $g^{hi}$  the metric matrix of  $U_qso(N)$ , which was defined in (5.7). The square roots of the diagonal elements  $\mathcal{L}_i^{-i}$ , as well as the square roots of the diagonal elements  $\mathcal{L}_i^{+i}$ , generate the same Cartan subalgebra of  $H$ , which coincides with  $H^+ \cap H^-$ . The braid matrix  $\hat{R}$  of  $U_qsl(N)$  is given by

$$\hat{R} = q^{-\frac{1}{N}} \left[ q \sum_i e_i^i \otimes e_i^i + \sum_{i \neq j} e_i^j \otimes e_j^i + k \sum_{i < j} e_i^i \otimes e_j^j \right] \quad (\text{A.9})$$

where all indices  $i, j, a, \dots = 1, 2, \dots, N$ , and  $e_j^i$  is the  $N \times N$  matrix with all elements equal to zero except for a 1 in the  $i$ th column and  $j$ th row. When  $H = U_qso(N)$  it is convenient to adopt the convention that all indices  $i, j, a, \dots$  take the values  $i = -n, \dots, -1, 0, 1, \dots, n$  for  $N$  odd, and  $i = -n, \dots, -1, 1, \dots, n$  for  $N$  even, where  $n := \lfloor \frac{N}{2} \rfloor$  is the rank of  $so(N)$ . Then the corresponding braid matrix reads

$$\begin{aligned} \hat{R} = & q \sum_{i \neq 0} e_i^i \otimes e_i^i + \sum_{\substack{i \neq j, -j \\ \text{or } i=j=0}} e_i^j \otimes e_j^i + q^{-1} \sum_{i \neq 0} e_i^{-i} \otimes e_{-i}^i \quad (\text{A.10}) \\ & + k \left( \sum_{i < j} e_i^i \otimes e_j^j - \sum_{i < j} q^{-\rho_i + \rho_j} e_i^{-j} \otimes e_{-i}^j \right). \end{aligned}$$

The braid matrix of  $sl(N)$  admits the orthogonal projector decomposition

$$q^{\frac{1}{N}} \hat{R} = q \mathcal{P}_S - q^{-1} \mathcal{P}_a, \quad \mathfrak{g} = sl(N); \quad (\text{A.11})$$

$\mathcal{P}_a, \mathcal{P}_S$  are the  $U_q sl(N)$ -covariant deformed antisymmetric and symmetric projectors. The braid matrix of  $so(N)$  admits the orthogonal projector decomposition

$$\hat{R} = q\mathcal{P}_s - q^{-1}\mathcal{P}_a + q^{1-N}\mathcal{P}_t \quad \mathbf{g} = so(N); \quad (\text{A.12})$$

$\mathcal{P}_a, \mathcal{P}_t, \mathcal{P}_s$  are the corresponding  $q$ -deformed antisymmetric, trace, trace-free symmetric projectors.

By iterated use of equations (A.5-A.7) one immediately shows that they hold also if  $\hat{R}$  is replaced by any polynomial function  $f(\hat{R})$  of  $\hat{R}$ , in particular by  $f(\hat{R}) = \hat{R}^{-1}, \mathcal{P}_a$ .

Finally, the coproduct of the FRT generators is given by

$$\Delta(\mathcal{L}^{+i}_j) = \mathcal{L}^{+i}_h \otimes \mathcal{L}^{+h}_j, \quad \Delta(\mathcal{L}^{-i}_j) = \mathcal{L}^{-i}_h \otimes \mathcal{L}^{-h}_j. \quad (\text{A.13})$$

When  $H = U_q so(N)$  the antipode is given by

$$S\mathcal{L}^{\mp j}_i = g_{ih}\mathcal{L}^{\mp h}_k g^{kj}. \quad (\text{A.14})$$

The non-compact real sections of  $U_q \mathbf{g}$  require  $|q| = 1$  and are characterized by the  $*$ -structure

$$(\mathcal{L}^{\pm i}_j)^* = U^{-1i}_r \mathcal{L}^{\pm r}_s U^s_j. \quad (\text{A.15})$$

Here one can take

$$U^i_j := \begin{cases} g^{ih}g_{jh} & \text{if } \mathbf{g} = so(N), \\ q^{-i}\delta^i_j & \text{if } \mathbf{g} = sl(N) \end{cases}. \quad (\text{A.16})$$

The compact real section of  $U_q \mathbf{g}$  requires  $q \in \mathbb{R}^+$  if  $\mathbf{g} = so(N)$ ,  $q \in \mathbb{R}$  if  $\mathbf{g} = sl(N)$  and is characterized by the  $*$ -structure

$$(\mathcal{L}^{\pm i}_j)^* = S\mathcal{L}^{\mp j}_i. \quad (\text{A.17})$$

For  $\mathbf{g} = so(N)$  this amounts to

$$(\mathcal{L}^{\pm i}_j)^* = g_{ih}\mathcal{L}^{\mp h}_k g^{kj}. \quad (\text{A.18})$$

## Proof of Lemma 1

We start by recalling two relations proved in Lemma 2 of Ref. [2]

$$\begin{aligned} \mu_a \tilde{\varphi}^-(S\mathcal{L}^{-i}_b) &= \hat{R}^{-1cd}_{ab} \tilde{\varphi}^-(S\mathcal{L}^{-i}_c) \mu_d, \\ \bar{\mu}_a \tilde{\varphi}^+(S\mathcal{L}^{+i}_b) &= \hat{R}^{cd}_{ab} \tilde{\varphi}^+(S\mathcal{L}^{+i}_c) \bar{\mu}_d. \end{aligned} \quad (\text{A.19})$$

Applying  $\tilde{\varphi}^{\pm}$  to (5.5) we find

$$\tilde{\varphi}^{\pm}(S\mathcal{L}^{\pm a}_b) p^i = \hat{R}^{\pm 1 ai}_{jk} p^j \tilde{\varphi}^{\pm}(S\mathcal{L}^{\pm k}_b). \quad (\text{A.20})$$

We finally note that

$$\bar{\mu}_{\pm 1} = -\mu_{\mp 1} \frac{p^{\mp 1}}{p^{\pm 1}} q^2. \quad (\text{A.21})$$

The claim is a direct consequence of relations (A.19), (A.20), (A.21).

## Proof of Proposition 10

For odd  $N$

$$\begin{aligned}
\text{lhs}(5.25) &\stackrel{(5.15)}{=} \sum_a \mathcal{L}_a^{+h} \tilde{\varphi}^+(S\mathcal{L}_j^{+a}) \zeta_5^-(\mathcal{L}_k^{-i}) \\
&\stackrel{\text{Thm 4}}{=} \sum_a \mathcal{L}_a^{+h} \zeta_5^-(\mathcal{L}_k^{-i}) \tilde{\varphi}^+(S\mathcal{L}_j^{+a}) \\
&\stackrel{(5.15)}{=} \sum_{a,b} \mathcal{L}_a^{+h} \mathcal{L}_b^{-i} \tilde{\varphi}^-(S\mathcal{L}_k^{-b}) \tilde{\varphi}^+(S\mathcal{L}_j^{+a}) \\
&\stackrel{(5.22)}{=} \sum_{a,b} \sum_{l,m,r,s} \mathcal{L}_a^{+h} \mathcal{L}_b^{-i} \hat{R}_{lm}^{-1ba} \tilde{\varphi}^+(S\mathcal{L}_r^{+l}) \tilde{\varphi}^-(S\mathcal{L}_s^{-m}) \hat{R}_{kj}^{rs} \frac{\gamma_k}{\bar{\gamma}_k} \frac{\bar{\gamma}_s}{\gamma_s} \\
&\stackrel{(A.7)}{=} \sum_{c,d} \sum_{l,m,r,s} \hat{R}_{cd}^{-1ih} \mathcal{L}_m^{-d} \mathcal{L}_l^{+c} \tilde{\varphi}^+(S\mathcal{L}_r^{+l}) \tilde{\varphi}^-(S\mathcal{L}_s^{-m}) \hat{R}_{kj}^{rs} \frac{\gamma_k}{\bar{\gamma}_k} \frac{\bar{\gamma}_s}{\gamma_s} \\
&\stackrel{(5.15)}{=} \sum_{c,d} \sum_{m,r,s} \hat{R}_{cd}^{-1ih} \mathcal{L}_m^{-d} \zeta_5^+(\mathcal{L}_r^{+c}) \tilde{\varphi}^-(S\mathcal{L}_s^{-m}) \hat{R}_{kj}^{rs} \frac{\gamma_k}{\bar{\gamma}_k} \frac{\bar{\gamma}_s}{\gamma_s} \\
&\stackrel{\text{Thm 4}}{=} \sum_{c,d} \sum_{m,r,s} \hat{R}_{cd}^{-1ih} \mathcal{L}_m^{-d} \tilde{\varphi}^-(S\mathcal{L}_s^{-m}) \zeta_5^+(\mathcal{L}_r^{+c}) \hat{R}_{kj}^{rs} \frac{\gamma_k}{\bar{\gamma}_k} \frac{\bar{\gamma}_s}{\gamma_s} \\
&\stackrel{(5.15)}{=} \text{rhs}(5.25).
\end{aligned}$$

For even  $N$

$$\begin{aligned}
&\zeta_5^+(\mathcal{L}_j^{+h}) \zeta_5^-(\mathcal{L}_k^{-i}) \stackrel{(5.15)}{=} \sum_b \mathcal{L}_b^{+h} \tilde{\varphi}^+(S\mathcal{L}_j^{+b}) \zeta_5^-(\mathcal{L}_k^{-i}) \\
&\stackrel{\text{Thm 4}, (5.19)}{=} \sum_b \mathcal{L}_b^{+h} \zeta_5^-(\mathcal{L}_k^{-i}) \tilde{\varphi}^+(S\mathcal{L}_j^{+b}) q^{\eta_j(\eta_i - \eta_k)} \\
&\stackrel{(5.15)}{=} \sum_{a,b} \mathcal{L}_b^{+h} \mathcal{L}_a^{-i} \tilde{\varphi}^-(S\mathcal{L}_k^{-a}) \varphi^+(S\mathcal{L}_j^{+b}) q^{\eta_j(\eta_i - \eta_k)} \\
&\stackrel{(A.7)}{=} \sum_{a,b} \sum_{l,m,c,d} \hat{R}_{lm}^{-1ih} \mathcal{L}_d^{-m} \mathcal{L}_c^{+l} \hat{R}_{ab}^{cd} \tilde{\varphi}^-(S\mathcal{L}_k^{-a}) \varphi^+(S\mathcal{L}_j^{+b}) q^{\eta_j(\eta_i - \eta_k)}.
\end{aligned}$$

If  $k \notin \{1, -1\}$  then

$$\begin{aligned}
&\zeta_5^+(\mathcal{L}_j^{+h}) \zeta_5^-(\mathcal{L}_k^{-i}) \stackrel{(5.23)}{=} \frac{\gamma_k}{\bar{\gamma}_k} \sum_{l,m,c,d} \hat{R}_{lm}^{-1ih} \mathcal{L}_d^{-m} \mathcal{L}_c^{+l} \left[ \sum_{\substack{r,s \\ s \neq \pm 1}} \tilde{\varphi}^+(S\mathcal{L}_r^{+c}) \right. \\
&\quad \left. \times \tilde{\varphi}^-(S\mathcal{L}_s^{-d}) \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} + \sum_{\substack{r,s \\ s = \pm 1}} \tilde{\varphi}^+(S\mathcal{L}_r^{+c}) \tilde{\varphi}^-(S\mathcal{L}_{-s}^{-d}) \frac{p^{-s}}{p^s} \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^2 \right] q^{\eta_j(\eta_i - \eta_k)} \\
&\stackrel{(5.15)}{=} \frac{\gamma_k}{\bar{\gamma}_k} \sum_{l,m,d,r} \hat{R}_{lm}^{-1ih} \mathcal{L}_d^{-m} \zeta_5^+(\mathcal{L}_r^{+l}) \left[ \sum_{s \neq \pm 1} \tilde{\varphi}^-(S\mathcal{L}_s^{-d}) \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} \right.
\end{aligned}$$



$$\begin{aligned}
& + \sum_{s=\pm 1} \tilde{\varphi}^-(S\mathcal{L}_{-s}^{-d}) \frac{p^{-s}}{p^s} \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^2 \Big] q^{\eta_j(\eta_i - \eta_k)} \\
& \stackrel{\text{Thm 4, (5.19)}}{=} \frac{\gamma_k}{\bar{\gamma}_k} \sum_{l,m,d,r} \hat{R}_{lm}^{-1ih} \mathcal{L}_d^{-m} \left[ \sum_{s \neq \pm 1} \tilde{\varphi}^-(S\mathcal{L}_s^{-d}) \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} \right. \\
& + \sum_{s=\pm 1} \tilde{\varphi}^-(S\mathcal{L}_{-s}^{-d}) \frac{p^{-s}}{p^s} \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^{2+\eta_s(\eta_r - \eta_l)} \Big] \zeta_5^+(\mathcal{L}_r^+ q^{\eta_j(\eta_i - \eta_k)} \\
& \stackrel{(5.15)}{=} \frac{\gamma_k}{\bar{\gamma}_k} \sum_{l,m,r} \hat{R}_{lm}^{-1ih} \left[ \sum_{s \neq \pm 1} \zeta_5^-(\mathcal{L}_s^{-m}) \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} + \right. \\
& \left. \sum_{s=\pm 1} \zeta_5^-(\mathcal{L}_{-s}^{-m}) \frac{p^{-s}}{p^s} \hat{R}_{kj}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^{2+\eta_s(\eta_r - \eta_l)} \right] \zeta_5^+(\mathcal{L}_r^+ q^{\eta_j(\eta_i - \eta_k)} \\
& \stackrel{\text{Thm 4}}{=} \text{rhs}(5.26).
\end{aligned}$$

If  $k \in \{1, -1\}$  then

$$\begin{aligned}
& \zeta_5^+(\mathcal{L}_j^+ \zeta_5^-(\mathcal{L}_k^-) \stackrel{(5.23)}{=} - \sum_{\substack{l,m,c \\ d,r}} \hat{R}_{lm}^{-1ih} \mathcal{L}_d^{-m} \mathcal{L}_c^+ \tilde{\varphi}^+(S\mathcal{L}_r^+) \left[ \sum_{s \neq \pm 1} \tilde{\varphi}^-(S\mathcal{L}_s^{-d}) \right. \\
& \times \frac{\bar{\gamma}_s}{\gamma_s} \hat{R}_{-k,j}^{rs} + \sum_{s=\pm 1} \tilde{\varphi}^-(S\mathcal{L}_{-s}^{-d}) \frac{p^{-s}}{p^s} \hat{R}_{-k,j}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^2 \Big] q^{\eta_j(\eta_i - \eta_k)} \frac{p^{-k}}{p^k} q^{-2+2\eta_j\eta_k} \\
& \stackrel{(5.15)}{=} - \sum_{l,m,d,r} \hat{R}_{lm}^{-1ih} \mathcal{L}_d^{-m} \zeta_5^+(\mathcal{L}_r^+) \left[ \sum_{s \neq \pm 1} \tilde{\varphi}^-(S\mathcal{L}_s^{-d}) \hat{R}_{-k,j}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} \right. \\
& + \sum_{s=\pm 1} \tilde{\varphi}^-(S\mathcal{L}_{-s}^{-d}) \frac{p^{-s}}{p^s} \hat{R}_{-k,j}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^2 \Big] q^{-2+\eta_j(\eta_i + \eta_k)} \frac{p^{-k}}{p^k} \\
& \stackrel{\text{Thm 4, (5.19)}}{=} - \sum_{l,m,r} \hat{R}_{lm}^{-1ih} \mathcal{L}_d^{-m} \left[ \sum_{s \neq \pm 1} \tilde{\varphi}^-(S\mathcal{L}_s^{-d}) \hat{R}_{-k,j}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} \right. \\
& + \sum_{s=\pm 1} \tilde{\varphi}^-(S\mathcal{L}_{-s}^{-d}) \frac{p^{-s}}{p^s} \hat{R}_{-k,j}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^{2+\eta_s(\eta_r - \eta_l)} \Big] \zeta_5^+(\mathcal{L}_r^+) q^{-2+\eta_j(\eta_i + \eta_k)} \frac{p^{-k}}{p^k} \\
& \stackrel{(5.15)}{=} - \sum_{l,m,r} \hat{R}_{lm}^{-1ih} \left[ \sum_{s \neq \pm 1} \zeta_5^-(\mathcal{L}_s^{-m}) \hat{R}_{-k,j}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} + \sum_{s=\pm 1} \zeta_5^-(\mathcal{L}_{-s}^{-m}) \right. \\
& \times \frac{p^{-s}}{p^s} \hat{R}_{-k,j}^{rs} \frac{\bar{\gamma}_s}{\gamma_s} q^{2+\eta_s(\eta_r - \eta_l)} \Big] \zeta_5^+(\mathcal{L}_r^+) q^{-2+\eta_j(\eta_i + \eta_k)} \frac{p^{-k}}{p^k} \\
& \stackrel{\text{Thm 4}}{=} \text{rhs}(5.27).
\end{aligned}$$

## References

- [1] U. Carow-Watamura, M. Schlieker, S. Watamura, “ $SO_q(N)$  covariant differential calculus on quantum space and quantum deformation of Schroedinger equation”, *Z. Physik C - Particles and Fields* **49** (1991), 439.
- [2] B. L. Cerchiai, G. Fiore, J. Madore, “Geometrical Tools for Quantum Euclidean Spaces”, *Commun. Math. Phys.* **217** (2001), 521-554.
- [3] B.L. Cerchiai, J. Madore, S. Schraml, J. Wess, “Structure of the Three-dimensional Quantum Euclidean Space”, *Eur. Phys. J. C* **16** (2000), 169.
- [4] C.-S. Chu, B. Zumino, “Realization of vector fields for quantum groups as pseudodifferential operators on quantum spaces”, Proc. XX Int. Conf. on Group Theory Methods in Physics, Toyonaka (Japan), 1995, and q-alg/9502005.
- [5] V. Drinfeld, “Quantum groups,” in *I.C.M. Proceedings, Berkeley*, p. 798. 1986.
- [6] L.D. Faddeev, N.Y. Reshetikhin, L. Takhtadjan, “Quantization of Lie groups and Lie algebras”, *Alge. i Anal.* **1** (1989), 178, translated from the Russian in *Leningrad Math. J.* **1** (1990), 193.
- [7] G. Fiore, “Realization of  $U_q(so(N))$  within the Differential Algebra on  $\mathbf{R}_q^N$ ”,
- [8] G. Fiore, “The Euclidean Hopf algebra  $U_q(e^N)$  and its fundamental Hilbert space representations”, *J. Math. Phys.* **36** (1995), 4363-4405; “The  $q$ -Euclidean Algebra  $U_q(e^N)$  and the Corresponding  $q$ -Euclidean Lattice”, *Int. J. Mod. Phys. A* **11** (1996), 863-886.
- [9] G. Fiore e J. Madore “The geometry of the quantum Euclidean space” *J. Geom. Phys.* **33** (2000), 257-287.
- [10] G. Fiore, H. Steinacker and J. Wess “Unbraiding the braided tensor product”, math/0007174.
- [11] O. Ogievetsky “Differential operators on quantum spaces for  $GL_q(n)$  and  $SO_q(n)$ ” *Lett. Math. Phys.* **24** (1992), 245.
- [12] H. Grosse, J. Madore, H. Steinacker, “Field Theory on the  $q$ -deformed Fuzzy Sphere”, *J. Geom. Phys.* **38** (2001), 308-342.
- [13] T. Hayashi, “ $q$ -Analogues of Clifford and Weyl Algebras: Spinor and Oscillator Realizations of Quantum Enveloping algebras” *Commun. Math. Phys.* **127** (1990), 129.
- [14] O. Ogievetsky, W.B. Schmidke, J. Wess, B. Zumino, “ $q$ -deformed Poincaré algebra” *Commun. Math. Phys.* **150** (1992), 495.
- [15] W. Pusz, S. L. Woronowicz, “Twisted Second Quantization”, *Rep. Math. Phys.* **27** (1989), 231.

- [16] J. Wess, B. Zumino, “Covariant differential calculus on the quantum hyperplane”, *Nucl. Phys. (Proc. Suppl.)* **18B** (1990) 302.